

ON DERIVED EQUIVALENCE CLASSIFICATION OF GENTLE TWO-CYCLE ALGEBRAS

GRZEGORZ BOBIŃSKI AND PIOTR MALICKI (TORUŃ)

ABSTRACT. We classify, up to derived (equivalently, tilting-cotilting) equivalence all nondegenerate gentle two-cycle algebras. We also give a partial classification and formulate a conjecture in the degenerate case.

INTRODUCTION AND THE MAIN RESULT

Throughout the paper k denotes a fixed algebraically closed field. By an algebra we mean a finite dimensional basic connected k -algebra and by a module a finite dimensional left module. By \mathbb{Z} , \mathbb{N} , and \mathbb{N}_+ , we denote the sets of integers, nonnegative integers, and positive integers, respectively. Finally, if $i, j \in \mathbb{Z}$, then $[i, j] = \{l \in \mathbb{Z} \mid i \leq l \leq j\}$.

With an algebra Λ we may associate its bounded derived category $\mathcal{D}^b(\Lambda)$ (in the sense of Verdier [29]) of bounded complexes of Λ -modules, which has a structure of a triangulated category (see [17]). The bounded derived category is an important homological invariant of the module category of an algebra and attracts a lot of interest (see for example [5, 8, 15, 16, 18, 22, 24, 25]). In particular, the derived equivalence classes of algebras have been investigated (see for example [1, 9, 11, 14, 20]), where two algebras are said to be derived equivalent if their bounded derived categories are equivalent as triangulated categories.

A handy way of proving a derived equivalence between algebras Λ and Λ' is a construction of a (co)tilting Λ -module T such that Λ' is (isomorphic to) the opposite of the endomorphism algebra of T . Here a Λ -module T is called (co)tilting if $\text{pd}_\Lambda T \leq 1$ ($\text{id}_\Lambda T \leq 1$, respectively), $\text{Ext}_\Lambda^1(T, T) = 0$, and T is a direct sum of precisely $\text{rk } K_0(\Lambda)$ pairwise nonisomorphic indecomposable Λ -modules, where $K_0(\Lambda)$ denotes the Grothendieck group of the category of Λ -modules. The transitive closure of the relation defined in this way is called tilting-cotilting equivalence. For many classes of algebras tilting-cotilting equivalence and derived equivalence coincide.

Results of this type have been obtained for gentle algebras, introduced by Assem and Skowroński in [4] (see Section 1 for a precise definition), which form an important subclass of the class of special

2000 *Mathematics Subject Classification.* 18E30, 16G20.


Key words and phrases. derived category, gentle algebra, tilting-cotilting equivalence.

biserial algebras in the sense of [27]. We note that a representation-infinite algebra admits a simply connected Galois covering whose every finite convex subcategory is representation-finite if and only if it is a special biserial algebra and its simply connected Galois covering is the repetitive category of union of a countable chain of gentle tree algebras (see [23]). The class of algebras derived equivalent to a hereditary algebra of Dynkin type \mathbb{A}_n for some $n \in \mathbb{N}_+$ coincides with the class of algebras tilting-cotilting equivalent to a hereditary algebra of type \mathbb{A}_n and consists of the gentle algebras whose Gabriel quivers have n vertices and $n-1$ arrows (see [2]). Moreover, for a given n all such algebras form one derived equivalence class. Similarly, the class of algebras derived equivalent to a hereditary algebra of Euclidean type $\tilde{\mathbb{A}}_n$ for some $n \in \mathbb{N}_+$ coincides with the class of algebras tilting-cotilting equivalent to a hereditary algebra of type $\tilde{\mathbb{A}}_n$ and consists of the gentle algebras whose Gabriel quivers have n vertices and n arrows and which satisfy the so-called clock condition on the unique cycle. In this case, there are exactly $\lfloor \frac{n}{2} \rfloor$ derived (equivalently, tilting-cotilting) equivalence classes for a given n . The algebras with the same numbers of vertices and arrows in the Gabriel quiver equal are called one-cycle algebras. The remaining gentle one-cycle algebras form the class of derived discrete algebras which are not derived (equivalently, tilting-cotilting) equivalent to a hereditary algebra of Dynkin type (see [30]). The derived equivalence classes of these algebras were described in [10].

The aim of this paper is to extend the above classification to the class of gentle two-cycle algebras, where we call an algebra a two-cycle algebra if the number of arrows in the Gabriel quiver exceeds the number of vertices by one. An additional motivation for this research is the fact proved by Schröer and Zimmermann in [26] saying that the gentle algebras are closed under derived equivalences. Moreover, for the gentle algebras the numbers of vertices and arrows in the Gabriel quiver are derived invariants (see [7, Corollary 15]). However, we obtain a full classification only for nondegenerate gentle two-cycle algebras, where we call a gentle two-cycle algebra Λ nondegenerate if $\sum_{(n,m) \in \mathbb{N} \times \mathbb{N}} \phi_\Lambda(n, m) = 3$ and $\phi_\Lambda : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ is the derived invariant introduced by Avella-Alaminos and Geiss in [7] (see Section 3). For the remaining gentle two-cycle algebras Λ , which we call degenerate, we have $\sum_{(n,m) \in \mathbb{N} \times \mathbb{N}} \phi_\Lambda(n, m) = 1$. Obviously, the both classes of gentle two-cycle algebras are closed under derived (hence also tilting-cotilting) equivalences.


Before formulating the main results of the paper we define the following families of algebras.

- $\Lambda_0(p, r)$ for $p \in \mathbb{N}_+$ and $r \in [0, p-1]$ is the algebra of the quiver




bound by $\alpha_p\beta$, $\alpha_i\alpha_{i+1}$ for $i \in [1, r]$ and $\gamma\alpha_1$.

- $\Lambda'_0(p, r)$ for $p \in \mathbb{N}_+$ and $r \in [0, p-1]$ is the algebra of the quiver




bound by $\alpha_i\alpha_{i+1}$ for $i \in [1, r]$, $\alpha_p\gamma$, and $\beta\delta$.

- $\Lambda_1(p_1, p_2, p_3, p_4, r_1)$ for $p_1, p_2 \in \mathbb{N}_+$, $p_3, p_4 \in \mathbb{N}$, and $r_1 \in [0, p_1-1]$, such that $p_2 + p_3 \geq 2$ and $p_4 + r_1 \geq 1$, is the algebra of the quiver



bound by $\alpha_i\alpha_{i+1}$ for $i \in [p_1 - r_1, p_1 - 1]$, $\alpha_{p_1}\beta_1$, $\beta_i\beta_{i+1}$ for $i \in [1, p_2 - 1]$, and $\beta_{p_2}\alpha_1$.

- $\Lambda_2(p_1, p_2, p_3, r_1, r_2)$ for $p_1, p_2 \in \mathbb{N}_+$, $p_3 \in \mathbb{N}$, $r_1 \in [0, p_1 - 1]$, and $r_2 \in [0, p_2 - 1]$, such that $p_3 + r_1 + r_2 \geq 1$, is the algebra of the quiver



bound by $\alpha_i\alpha_{i+1}$ for $i \in [p_1 - r_1, p_1 - 1]$, $\alpha_{p_1}\alpha_1$, $\beta_i\beta_{i+1}$ for $i \in [p_2 - r_2, p_2 - 1]$, and $\beta_{p_2}\beta_1$.

The main results of the paper are the following.

Theorem 1. *If Λ is a nondegenerate gentle two-cycle algebra, then Λ is derived (equivalently, tilting-cotilting) equivalent to one of the following algebras:*

- $\Lambda_1(p_1, p_2, p_3, p_4, r_1)$ for some $p_1, p_2 \in \mathbb{N}_+$, $p_3, p_4 \in \mathbb{N}$, and $r_1 \in [0, p_1 - 1]$, such that $p_2 + p_3 \geq 2$, $p_4 + r_1 \geq 1$, and either $p_3 > p_4$ or $p_3 = p_4$ and $p_2 > r_1$,
- $\Lambda_2(p_1, p_2, p_3, r_1, r_2)$ for some $p_1, p_2 \in \mathbb{N}_+$, $p_3 \in \mathbb{N}$, $r_1 \in [0, p_1 - 1]$, and $r_2 \in [0, p_2 - 1]$, such that $p_3 + r_1 + r_2 \geq 1$ and either $p_1 > p_2$ or $p_1 = p_2$ and $r_1 \geq r_2$.

Moreover, different algebras from the above list are not derived (equivalently, tilting-cotilting) equivalent.

Theorem 2. *If Λ is a degenerate gentle two-cycle algebra, then Λ is derived (equivalently, tilting-cotilting) equivalent to one of the following algebras:*

- $\Lambda_0(p, r)$ for some $p \in \mathbb{N}_+$ and $r \in [0, p - 1]$,
- $\Lambda'_0(p, 0)$ for some $p \in \mathbb{N}_+$.

Moreover, we have the following conjecture concerning the minimality of the list in the above theorem.

Conjecture. *Different algebras from the list in Theorem 2 are not derived (equivalently, tilting-cotilting) equivalent.*

Obviously, if $p_1, p_2 \in \mathbb{N}_+$, $r_1 \in [0, p_1 - 1]$, $r_2 \in [0, p_2 - 1]$, and $p_1 \neq p_2$, then $\Lambda_0(p_1, r_1)$ and $\Lambda_0(p_2, r_2)$ ($\Lambda'_0(p_1, 0)$ and $\Lambda'_0(p_2, 0)$, respectively) are not derived equivalent. Similarly, if $p_1, p_2 \in \mathbb{N}_+$, $r_1 \in [0, p_1 - 1]$, and $p_1 \neq p_2 + 1$, then $\Lambda_0(p_1, r_1)$ and $\Lambda'_0(p_2, 0)$ are not derived equivalent. Thus it is enough to prove that $\Lambda_0(p+1, 0), \dots, \Lambda_0(p+1, p)$ and $\Lambda'_0(p, 0)$ are pairwise not derived equivalent for a fixed $p \in \mathbb{N}_+$. It follows easily by investigating the Euler quadratic forms that $\Lambda_0(p+1, r_1)$ and $\Lambda(p+1, r_2)$ ($\Lambda'_0(p, 0)$ and $\Lambda_0(p+1, r_2)$) are not derived equivalent if $r_1 \not\equiv r_2 \pmod{2}$ ($r_2 \equiv 0 \pmod{2}$), respectively).

The paper is organized as follows. In Section 1 we first present basic definitions, then describe main tools used in order to reduce an arbitrary gentle two-cycle algebra to one of the algebras listed in Theorems 1 and 2: passing to the opposite algebra, (generalized) APR-(co)reflections, and HW-(co)reflections. Finally, we describe in Section 1 an operation of shifting relations being a basic application of the above operations, and investigate two particular families of gentle two-cycle algebras. In Section 2, being a technical heart of the paper, we prove, in a sequence of steps, that the lists of representatives of the tilting-cotilting equivalence classes of gentle two-cycle algebras given in Theorems 1 and 2 are complete, while in Section 3 we show that different algebras from the list given in Theorem 1 are not derived equivalent. The last property follows from calculations of the derived invariant introduced by Avella-Alaminos and Geiss in [7].

For a basic background on representation theory of finite dimensional algebras we refer to [3].

Authors gratefully acknowledges the support from the Polish Scientific Grant KBN No. 1 P03A 018 27.

1. BASIC TOOLS AND AUXILIARY RESULTS

By a (finite) quiver Δ we mean a finite set Δ_0 of vertices together with a finite set Δ_1 of arrows and two maps $s = s_\Delta, t = t_\Delta : \Delta_1 \rightarrow \Delta_0$ which assign to an arrow α its starting and terminating vertex, respectively. We say that an arrow α is adjacent to a vertex x if either $s\alpha = x$ or $t\alpha = x$. By a path of length $n \in \mathbb{N}_+$ we mean a sequence $\sigma = \alpha_1 \cdots \alpha_n$ of arrows such that $s\alpha_i = t\alpha_{i+1}$ for all $i \in [1, n-1]$. In the above situation we denote $s\alpha_n$ and $t\alpha_1$ by $s\sigma$ and $t\sigma$, respectively. We also call α_1 and α_n the terminating and the starting arrow of σ , respectively. Additionally, for each $x \in \Delta_0$ we consider the trivial path of length 0, also denoted by x , such that $sx = x = tx$. The length of a path σ will be denoted by $\ell(\sigma)$. A path σ is called maximal if there exists no arrow α such that either $s\alpha = t\sigma$ or $t\alpha = s\sigma$. Similarly, we define maximal paths starting (or terminating) at a given vertex. A connected quiver is said to be c -cycle if $|\Delta_1| = |\Delta_0| + c - 1$.

With a quiver Δ we associate its path algebra $k\Delta$, which as a k -vector space has a basis formed by all paths in Δ and whose multiplication is induced by the composition of paths. By a relation ρ in Δ we mean a linear combination of paths of length at least 2 with common starting and terminating vertices. The common starting vertex is denoted by $s\rho$ and the common terminating vertex by $t\rho$. A set R of relations is called minimal if ρ does not belong to the ideal $\langle R \setminus \{\rho\} \rangle$ of $k\Delta$ generated by $R \setminus \{\rho\}$ for every $\rho \in R$. A pair (Δ, R) consisting of a quiver Δ and a minimal set of relations R such that there exists $n \in \mathbb{N}$ with $\sigma \in \langle R \rangle$ for each path σ in Δ of length at least n , is called a bound quiver. If (Δ, R) is a bound quiver, then the algebra $k\Delta/\langle R \rangle$ is called the bound quiver algebra of (Δ, R) .

Let (Δ, R) be a bound quiver and assume that R consists of paths. A path σ in Δ is said to be a path (Δ, R) if $\sigma \notin \langle R \rangle$ (in other words, none of the paths from R is a subpath of σ). A path σ in (Δ, R) is said to be maximal if there is no $\alpha \in \Delta_1$ such that either $s\alpha = t\sigma$ and $\alpha\sigma \notin \langle R \rangle$ or $t\alpha = s\sigma$ and $\sigma\alpha \notin \langle R \rangle$. Again we define maximal paths starting and terminating at a given vertex. If additionally R consists of paths of length two, then we say that $\alpha \in \Delta_1$ is a free arrow provided there exists no $\beta \in \Delta_1$ such that either $s\beta = t\alpha$ and $\beta\alpha \in R$ or $t\beta = s\alpha$ and $\alpha\beta \in R$.

Following [4] we say that a connected bound quiver (Δ, R) is gentle if the following conditions are satisfied:

- (1) for each $x \in \Delta_0$ there are at most two arrows α such that $s\alpha = x$ ($t\alpha = x$),
- (2) R consists of paths of length two,

- (3) for each $\alpha \in \Delta_1$ there is at most one arrow β such that $t\beta = s\alpha$ and $\alpha\beta \notin R$ ($s\beta = t\alpha$ and $\beta\alpha \notin R$),
- (4) for each $\alpha \in \Delta_1$ there is at most one arrow β such that $t\beta = s\alpha$ and $\alpha\beta \in R$ ($s\beta = t\alpha$ and $\beta\alpha \in R$).

An algebra which is isomorphic to the bound quiver algebra a gentle bound quiver is called gentle.

With an abelian category \mathcal{A} we may associate its bounded derived category $\mathcal{D}^b(\mathcal{A})$ in the following way (see for example [29] for details). The objects of $\mathcal{D}^b(\mathcal{A})$ are the bounded complexes of objects of \mathcal{A} and the morphisms are obtained from the morphisms in the homotopy category by formally inverting the quasi-isomorphisms (more precisely, by localizing with respect to the quasi-isomorphism), where by a quasi-isomorphism we mean a morphism of complexes which induces an isomorphism of homology groups. The derived category together with the shift functor sending X to the shifted complex $X[1]$, where $X[1]_n = X_{n+1}$ and $d_{X[1]}^n = -d_{X[1]}^{n+1}$ for $n \in \mathbb{Z}$, is a triangulated category (see for example [17]). We say that two abelian categories \mathcal{A} and \mathcal{B} are derived equivalent if there exists a triangle equivalence $\mathcal{D}^b(\mathcal{A}) \rightarrow \mathcal{D}^b(\mathcal{B})$. We say that two algebras Λ and Λ' (bound quivers (Δ, R) and (Δ', R')) are derived equivalent if their categories of modules (representations, respectively) are derived equivalent. It follows from [26, Corollary 1.2] and [7, Corollary 15] that for $c \in \mathbb{Z}$ gentle c -cycle algebras (bound quivers) are closed under derived equivalences.

Recall from [12, 19] that if Λ is an algebra, then a Λ -module T is called tilting if $\text{pd}_\Lambda T \leq 1$, $\text{Ext}_\Lambda^1(T, T) = 0$, and T is a direct sum of n pairwise nonisomorphic indecomposable modules, where n is the rank of the Grothendieck group of Λ . Dually, we define notion of a cotilting module. Algebras Λ and Λ' are said to be tilting-cotilting equivalent if there exists a sequence $\Lambda = \Lambda_0, \Lambda_1, \dots, \Lambda_n = \Lambda'$ of algebras such that for each $i \in [0, n-1]$ there exists a (co)tilting Λ_{i-1} -module T_{i-1} such that $\Lambda_i \simeq \text{End}_{\Lambda_{i-1}}(T_{i-1})^{\text{op}}$. It was proved by Happel [16, Corollary 1.7] that if Λ and Λ' are tilting-cotilting equivalent, then they are derived equivalent.

A vertex x in a quiver Δ is called a sink (source) if there is no $\alpha \in \Delta_1$ with $s\alpha = x$ ($t\alpha = x$, respectively). If x is a sink in a gentle bound quiver (Δ, R) , then we define a new gentle bound quiver (Δ', R') , called the bound quiver obtained from (Δ, R) by applying the APR-reflection at x , in the following way: $\Delta'_0 = \Delta_0$, $\Delta'_1 = \Delta_1$,

$$s_{\Delta'}\alpha = \begin{cases} x & \text{if } t_\Delta\alpha = x, \\ s_\Delta\alpha & \text{otherwise,} \end{cases}$$

$$t_{\Delta'}\alpha = \begin{cases} s_\Delta\alpha & \text{if } t_\Delta\alpha = x, \\ x & \text{if } \exists \beta \in \Delta_1 : t_\Delta\beta = x \wedge s_\Delta\beta = t_\Delta\alpha \wedge \beta\alpha \in R, \\ t_\Delta\alpha & \text{otherwise,} \end{cases}$$

and

$$R' = \{\rho \in R \mid t_\Delta \rho \neq x\} \cup \{\alpha\beta \mid t_\Delta \alpha = x \\ \wedge \exists \gamma \in \Delta_1 : \gamma \neq \alpha \wedge t_\Delta \gamma = x \wedge s_\Delta \gamma = t_\Delta \beta \wedge \gamma\beta \in R\}.$$

It follows that the bound quiver algebra of (Δ', R') is isomorphic to the opposite algebra of the endomorphism algebra of the APR-tilting module (see [6]) at x defined as

$$\bigoplus_{\substack{a \in \Delta_0 \\ a \neq x}} P(a) \oplus \left(\bigoplus_{\substack{\alpha \in \Delta_1 \\ t\alpha = x}} P(s\alpha) \right) / P(x)$$

(see [4, 2.1]).

We present now a generalization of the above construction due to Brenner and Butler (see [13, Chapter 2]). Let x be a vertex in a gentle bound quiver (Δ, R) such that there is no $\alpha \in \Delta_1$ with $s\alpha = x = t\alpha$ and for each $\alpha \in \Delta_1$ with $s\alpha = x$ there exists $\beta_\alpha \in \Delta_1$ with $t\beta_\alpha = x$ and $\alpha\beta_\alpha \notin R$. We define a bound quiver (Δ', R') in the following way: $\Delta'_0 = \Delta_0$, $\Delta'_1 = \Delta_1$,

$$s_{\Delta'} \alpha = \begin{cases} x & \text{if } t_\Delta \alpha = x, \\ s\beta_\alpha & \text{if } s_\Delta \alpha = x, \\ s_\Delta \alpha & \text{otherwise,} \end{cases}$$

$$t_{\Delta'} \alpha = \begin{cases} s_\Delta \alpha & \text{if } t_\Delta \alpha = x, \\ x & \text{if } \exists \beta \in \Delta_1 : t_\Delta \beta = x \wedge s_\Delta \beta = t_\Delta \alpha \wedge \beta\alpha \in R, \\ t_\Delta \alpha & \text{otherwise,} \end{cases}$$

and set

$$R' = \{\alpha\beta \in R \mid t_\Delta \alpha \neq x \wedge s_\Delta \alpha \neq x\} \cup \{\alpha\beta_\alpha \mid s_\Delta \alpha = x\} \cup \\ \{\alpha\beta \mid t_\Delta \alpha = x \wedge \exists \gamma \in \Delta_1 : \gamma \neq \alpha \wedge t_\Delta \gamma = x \wedge s_\Delta \gamma = t_\Delta \beta \wedge \gamma\beta \in R\}.$$

We will say that (Δ', R') is obtained from (Δ, R) by applying the generalized APR-reflection at x . Similarly as in the previous situation it follows easily that the bound quiver algebra of (Δ', R') is the opposite algebra of the endomorphism algebra of the tilting module defined in the same way as before. Obviously all APR-reflections are examples of generalized APR-reflections.

We also have a version of the above construction for a vertex x of a gentle bound quiver (Δ, R) such that there exists $\alpha \in \Delta_1$ with $s\alpha = x = t\alpha$. Observe that then $\alpha^2 \in R$. We additionally assume that there exists $\beta_0 \in \Delta_1$ with $s\beta_0 \neq x$ and $t\beta_0 = x$. We define a bound quiver (Δ', R') in the following way: $\Delta'_0 = \Delta_0$, $\Delta'_1 = \Delta_1$,

$$s_{\Delta'} \alpha = \begin{cases} x & \text{if } t_\Delta \alpha = x, \\ s_\Delta \beta_0 & \text{if } s_\Delta \alpha = x \wedge t_\Delta \alpha \neq x, \\ s_\Delta \alpha & \text{otherwise,} \end{cases}$$

$$t_{\Delta'}\alpha = \begin{cases} s_{\Delta}\alpha & \text{if } t_{\Delta}\alpha = x, \\ x & \text{if } \exists \beta \in \Delta_1 : t_{\Delta}\beta = x \wedge s_{\Delta}\beta = t_{\Delta}\alpha \wedge \beta\alpha \in R, \\ t_{\Delta}\alpha & \text{otherwise,} \end{cases}$$

and $R' = R$. We will say again that (Δ', R') is obtained from (Δ, R) by applying the generalized APR-reflection at x . It follows that the bound quiver algebra of (Δ', R') is the opposite algebra of the endomorphism algebra of the tilting module

$$\bigoplus_{\substack{a \in \Delta_0 \\ a \neq x}} P(a) \oplus (P(y) \oplus P(y))/P(x),$$

where $y = s\beta_0$ and $P(x)$ is embedded in $P(y) \oplus P(y)$ in such a way that the quotient module is indecomposable.

Let again x be a sink in a gentle bound quiver (Δ, R) . We define the HW-reflection of (Δ, R) at x as the bound quiver (Δ', R') constructed in the following way. If $\Delta_0 = \{x\}$ (equivalently, $\Delta_1 = \emptyset$), then $(\Delta', R') = (\Delta, R)$, hence assume this is not the case. Then we put $\Delta'_0 = \Delta_0$ and $\Delta'_1 = \Delta_1$. For each arrow α terminating at x let β_α be the starting arrow of the maximal path in (Δ, R) terminating at x whose terminating arrow is α . We put

$$s_{\Delta'}\alpha = \begin{cases} x & \text{if } t_{\Delta}\alpha = x, \\ s_{\Delta}\alpha & \text{otherwise,} \end{cases} \quad \text{and} \quad t_{\Delta'}\alpha = \begin{cases} s_{\Delta}\beta_\alpha & \text{if } t_{\Delta}\alpha = x, \\ t_{\Delta}\alpha & \text{otherwise.} \end{cases}$$

Finally let

$$R' = \{\rho \in R \mid t_{\Delta}\rho \neq x\} \cup \{\beta_\alpha \mid t_{\Delta}\alpha = x \wedge s_{\Delta}\beta = s_{\Delta}\beta_\alpha \wedge \beta \neq \beta_\alpha \wedge t_{\Delta}\beta \neq x\}.$$

It is known that bound quiver algebra of (Δ', R') is (isomorphic to) the algebra obtained from the bound quiver algebra of (Δ, R) by the HW-reflection at x (defined in [21]), hence in particular is tilting-cotilting equivalent to (Δ, R) (see [28]). Dually, one defines a quiver obtained from (Δ, R) by applying the HW-coreflection at a source.

Before we present basic applications of the above transformations, we describe one more construction. Let Σ be a subquiver of a quiver Δ . Assume that Σ' is a quiver such that $\Sigma'_0 = \Sigma_0$ and $\Sigma'_1 = \Sigma_1$ (but, usually, $s_{\Sigma'} \neq s_{\Sigma}$ and $t_{\Sigma'} \neq t_{\Sigma}$). We say that a quiver Δ' is obtained from Δ by replacing Σ by Σ' if $\Delta'_0 = \Delta_0$, $\Delta'_1 = \Delta_1$, and

$$s_{\Delta'}\alpha = \begin{cases} s_{\Delta}\alpha & \text{if } \alpha \in \Delta_1 \setminus \Sigma_1, \\ s_{\Sigma'}\alpha & \text{if } \alpha \in \Sigma_1, \end{cases} \quad t_{\Delta'}\alpha = \begin{cases} t_{\Delta}\alpha & \text{if } \alpha \in \Delta_1 \setminus \Sigma_1, \\ t_{\Sigma'}\alpha & \text{if } \alpha \in \Sigma_1, \end{cases}$$

for $\alpha \in \Delta_1$.

We describe now operations of shifting relations.

Lemma 1.1. *If*

$$\Sigma = \bullet_u \xleftarrow{\alpha_1} \bullet_x \xleftarrow{\alpha_2} \bullet_y \xleftarrow{\alpha_3} \bullet_v$$

is a subquiver of a gentle bound quiver (Δ, R) such that $\alpha_1\alpha_2 \in R$, $\alpha_2\alpha_3 \notin R$, and there are no other arrows adjacent to y , then (Δ, R) is tilting-cotilting equivalent to the bound quiver (Δ', R') , where $R' = R \setminus \{\alpha_1\alpha_2\} \cup \{\alpha_2\alpha_3\}$ and Δ' is obtained from Δ by replacing Σ by the quiver

$$\bullet_u \xleftarrow{\alpha_1} \bullet_y \xleftarrow{\alpha_2} \bullet_x \xleftarrow{\alpha_3} \bullet_z.$$

Proof. Apply the generalized APR-coreflection at y . \square

We remark that it may happen that one of the following equalities hold: $u = y$, $x = v$ or $u = v$ holds. Moreover, $u = y$ if and only if $x = v$, and in this case $\alpha_1 = \alpha_3$. We call the above operation shifting the relation $\alpha_1\alpha_2$ to the right. Dually, one defines the operation of shifting relations to the left.

We will need the following generalization of the above lemma.

Lemma 1.2. *If*

$$\Sigma = \bullet_u \xleftarrow{\alpha_1} \bullet_x \xleftarrow{\alpha_2} \bullet_{y_n} \xrightarrow{\beta_n} \bullet_{y_{n-1}} \cdots \xrightarrow{\beta_1} \bullet_{y_1} \xrightarrow{\alpha_3} \bullet_{y_0} \xleftarrow{\alpha_3} \bullet_v, \quad n \in \mathbb{N}_+,$$

is a subquiver of a gentle bound quiver (Δ, R) such that $\alpha_1\alpha_2 \in R$, β_1, \dots, β_n are free arrows, and there are no other arrows adjacent to x, y_0, \dots, y_n , then (Δ, R) is tilting-cotilting equivalent to the bound quiver (Δ', R') , where $R' = R \setminus \{\alpha_1\alpha_2\} \cup \{\alpha_2\alpha_3\}$ and Δ' is obtained from Δ by replacing Σ by the quiver

$$\bullet_u \xleftarrow{\alpha_1} \bullet_{y_0} \xrightarrow{\beta_1} \bullet_{y_1} \cdots \xrightarrow{\beta_n} \bullet_{y_{n-1}} \xrightarrow{\alpha_2} \bullet_{y_n} \xleftarrow{\alpha_3} \bullet_x \xleftarrow{\alpha_3} \bullet_z.$$

Proof. We leave it to the reader to verify that the following sequence of operations leads from (Δ, R) to (Δ', R') : first for each $i = n, \dots, 1$ we apply the APR-coreflections at y_i, \dots, y_n, x , and next we apply the generalized APR-coreflections at y_0, \dots, y_n . \square

We will also shift a group of relations in the following sense.

Lemma 1.3. *Let*

$$\Sigma = \bullet_y \xrightarrow{\beta} \bullet_{x_0} \xleftarrow{\alpha_1} \bullet_{x_1} \cdots \xleftarrow{\alpha_n} \bullet_{x_{n-1}} \xleftarrow{\alpha_n} \bullet_{x_n}, \quad n \geq 2,$$

be a subquiver of a gentle bound quiver (Δ, R) such that β is a free arrow, $\alpha_i\alpha_{i+1} \in R$ for all $i \in [1, n-1]$, and there are no other arrows adjacent to x_0, \dots, x_{n-1} . If there is no $\alpha \in \Delta_1$ with $t\alpha = x_n$ and $\alpha_n\alpha \in R$, then (Δ, R) is tilting-cotilting equivalent to the bound quiver

(Δ', R') , where $R' = R \setminus \{\alpha_{n-1}\alpha_n\} \cup \{\beta\alpha_1\}$ and Δ' is obtained from Δ by replacing Σ by the quiver

$$\bullet \xleftarrow{\beta} x_1 \xleftarrow{\alpha_1} \bullet \xleftarrow{\alpha_2} \bullet \cdots \xleftarrow{\alpha_{n-1}} x_{n-1} \xleftarrow{\alpha_n} \bullet \xrightarrow{\alpha_n} x_n \bullet.$$

Proof. We apply the APR-reflection at x_0 , followed by the composition of the APR-reflection at x_i and the generalized APR-reflection at x_0 applied for $i = 1, \dots, n-1$. \square

Observe that in the above lemma we shift relations to the left. Dually we define an operation of shifting a group of relations to the right.

We present now a reduction, being a consequence of the above lemmas, which will appear a few times in our proofs. Let

$$\Sigma = \bullet \xrightarrow{\alpha_1} x_1 \bullet \cdots \bullet \xrightarrow{\alpha_n} x_n \bullet, \quad n \in \mathbb{N}_+,$$

be a subquiver of a gentle bound quiver (Δ, R) such that there are no other arrows adjacent to x_1, \dots, x_{n-1} (it may happen that $x_0 = x_n$). We divide Σ_1 into two disjoint subsets $\Sigma_{1,+}$ and $\Sigma_{1,-}$ in such a way that, for each $i \in [1, n-1]$, α_i and α_{i+1} belong to the same subset if and only if either $s\alpha_i = t\alpha_{i+1}$ or $t\alpha_i = s\alpha_{i+1}$. We additionally assume that there exists $\varepsilon \in \{-, +\}$ such that $\alpha\beta \notin R$ for all $\alpha, \beta \in \Sigma_{1,\varepsilon}$ with $s\alpha = t\beta$. If $x_0 = t\alpha_1$, then by applying APR-reflections and shifts of relations (we leave details to the reader), hence by passing to a tilting-cotilting equivalent bound quiver, we may replace Σ by the quiver

$$\bullet \xleftarrow{\alpha'_1} \cdots \xleftarrow{\alpha'_{l_1}} \bullet \xrightarrow{\alpha''_{l_2}} \cdots \xrightarrow{\alpha''_1} \bullet \xleftarrow{\alpha'''_1} \cdots \xleftarrow{\alpha'''_{l_3}} \bullet$$

for some $l_1, l_2, l_3 \in \mathbb{N}$ with $l_1 + l_2 + l_3 = n$. Moreover, we may additionally assume that $l_3 = 0$ if either $x_n = t\alpha_n$ or $x_n = s\alpha_n$ and there is no $\alpha \in \Delta_1$ with $t\alpha = x_n$ and $\alpha_n\alpha \in R$. Obviously, we have the dual statement if $x_0 = s\alpha_1$.


The next observation is the following.

Lemma 1.4. *If $p_1, p_2 \in \mathbb{N}_+$, $p_3, p_4 \in \mathbb{N}$, and $r_1 \in [0, p_1 - 1]$, are such that $p_2 + p_3 \geq 2$ and $p_4 + r_1 \geq 1$, then $\Lambda_1(p_1, p_2, p_3, p_4, r_1)$ and $\Lambda_1(p_1 + p_2 - r_1 - 1, r_1 + 1, p_4, p_3, p_2 - 1)$ are tilting-cotilting equivalent.*

Proof. It follows immediately by shifting relations. \square

In order to formulate the next lemma we introduce a new family of algebras. Namely, for $p_1, p_2 \in \mathbb{N}_+$, $p_3, p_4 \in \mathbb{N}$, $r_1 \in [0, p_1 - 1]$, and $r_2 \in [0, p_2 - 1]$, such that $p_3 + p_4 + r_1 + r_2 \geq 1$, let $\Lambda'_2(p_1, p_2, p_3, p_4, r_1, r_2)$

be the algebra of the quiver



bound by $\alpha_i \alpha_{i+1}$ for $i \in [p_1 - r_1, p_1 - 1]$, $\alpha_{p_1} \alpha_1$, $\beta_i \beta_{i+1}$ for $i \in [p_2 - r_2, p_2 - 1]$, and $\beta_{p_2} \beta_1$.

Lemma 1.5. *If $p_1, p_2, p_3 \in \mathbb{N}_+$, $p_4 \in \mathbb{N}$, $r_1 \in [0, p_1 - 1]$, and $r_2 \in [0, p_2 - 1]$, then $\Lambda'_2(p_1, p_2, p_3, p_4, r_1, r_2)$ and $\Lambda'_2(p_1, p_2, p_3 - 1, p_4 + 1, r_1, r_2)$ are tilting-cotilting equivalent.*

Proof. Put $a_i = s\delta_i$, $i \in [1, p_4]$, and $b_i = s\beta_i$, $i \in [1, p_2]$. We first apply the APR-coreflections at a_{p_4}, \dots, a_1 , followed by the generalized APR-coreflection at b_{p_2} (we only apply the generalized APR-coreflection at b_{p_2} if $p_4 = 0$). Next we apply the APR-coreflection at b_{p_2-i} followed by the generalized APR-coreflection at b_{p_2} for $i = 1, \dots, r_2$ (we do nothing in this step if $r_2 = 0$, hence in particular we do nothing in this step if $p_2 = 1$), and finally we apply the APR-coreflections at $b_{p_2-r_2-1}, \dots, b_1$ (there is nothing to do if $r_2 = p_2 - 1$, hence again there is nothing to do if $p_2 = 1$). \square

Corollary 1.6. *If $p_1, p_2 \in \mathbb{N}_+$, $p_3 \in \mathbb{N}$, $r_1 \in [0, p_1 - 1]$, and $r_2 \in [0, p_2 - 1]$, are such that $p_3 + r_1 + r_2 \geq 1$, then $\Lambda_2(p_1, p_2, p_3, r_1, r_2)$ and $\Lambda_2(p_2, p_1, p_3, r_2, r_1)$ are tilting-cotilting equivalent.*

Proof. It follows immediately from the above lemma, since it is easily seen that $\Lambda_2(p_1, p_2, p_3, r_1, r_2)$ and $\Lambda_2(p_2, p_1, p_3, r_2, r_1)$ are isomorphic to $\Lambda'_2(p_1, p_2, p_3, 0, r_1, r_2)$ and $\Lambda'_2(p_1, p_2, 0, p_3, r_1, r_2)$, respectively. \square

Proposition 1.7. *If Λ is one of the algebras listed in Theorems 1 and 2, then Λ and Λ^{op} are tilting-cotilting equivalent.*


Proof. If either $\Lambda = \Lambda_0(p, r)$ for some $p \in \mathbb{N}_+$ and $r \in [0, p - 1]$ or $\Lambda = \Lambda'_0(p, 0)$ for some $p \in \mathbb{N}_+$, then the claim follows immediately by shifting relations. If $\Lambda = \Lambda_1(p_1, p_2, p_3, p_4, r_1)$ for some $p_1, p_2 \in \mathbb{N}_+$, $p_3, p_4 \in \mathbb{N}$, and $r_1 \in [0, p_1]$, such that $p_2 + p_3 \geq 2$ and $r_1 + p_4 \geq 1$, then we have to additionally apply APR-coreflections. Finally, if $\Lambda = \Lambda_2(p_1, p_2, p_3, r_1, r_2)$ for some $p_1, p_2 \in \mathbb{N}_+$, $p_3 \in \mathbb{N}$, $r_1 \in [0, p_1 - 1]$, and $r_2 \in [0, p_2 - 1]$, such that $p_3 + r_1 + r_2 \geq 1$, then the claim follows either from Corollary 1.6. \square

An important consequence of the above lemma is that in our considerations we may always replace an algebra by its opposite algebra. Indeed, if for an algebra Γ we are able to prove that Γ^{op} is tilting-cotilting equivalent to an algebra Λ listed in Theorems 1 and 2, then

obviously Γ is tilting-cotilting equivalent to Λ^{op} , hence also to Λ . In particular, once Theorems 1 and 2 are proved, then we know that if Γ is a gentle two-cycle algebra, then Γ and Γ^{op} are tilting-cotilting equivalent.

We finish this section by analyzing two particular families of gentle two-cycle bound quivers. First, we prove the following.

Proposition 1.8. *If (Δ, R) is a gentle bound quiver such that*



for some $p_1, p_2, q_1, q_2 \in \mathbb{N}_+$, then the bound quiver algebra of (Δ, R) is tilting-cotilting equivalent to $\Lambda'_0(p, r)$ for some $p \in \mathbb{N}_+$ and $r \in [0, p-1]$.

We first show that also in the proof of this theorem we may pass to the opposite algebras.

Lemma 1.9. *If $p \in \mathbb{N}_+$ and $r \in [1, p-1]$, then $\Lambda'_0(p, r)$ and $\Lambda_0(p+1, r-1)$ are tilting-cotilting equivalent.*


Proof. In order to prove this equivalence, we put $x = t\beta$, $z = s\delta$ and $x_1 = s\alpha_1$, and apply the APR-reflection at x followed by the APR-coreflection at z and the APR-reflection at x_1 to $\Lambda'_0(p, r)$. Then the claim follows by shifting relations. \square

Corollary 1.10. *If $p \in \mathbb{N}_+$ and $r \in [0, p-1]$, then $\Lambda'_0(p, r)$ and $\Lambda'_0(p, r)^{\text{op}}$ are tilting-cotilting equivalent.*

Proof. It follows either from Proposition 1.7 (if $r = 0$) or from the previous lemma and Proposition 1.7 (if $r > 0$). \square


In the proof of Proposition 1.8 we will need the following families of algebras:

- $\Gamma_0(p, q, r)$ for $p, q \in \mathbb{N}_+$ and $r \in [0, p-1]$ is the algebra of the quiver




bound by $\alpha_i\alpha_{i+1}$ for $i \in [p-r, p]$ and $\beta_q\beta_{q+1}$,

- $\Gamma_1(p, q, r, r')$ for $p, q \in \mathbb{N}_+$, $r \in [0, p - 1]$, and $r' \in \mathbb{N}$, is the algebra of the quiver



bound by $\alpha_i \alpha_{i+1}$ for $i \in [p - r, p + r']$ and $\beta_q \beta_{q+1}$,

- $\Gamma_2(p, q, r, r')$ for $p, q \in \mathbb{N}_+$, $r \in [0, p - 1]$, and $r' \in \mathbb{N}$, is the algebra of the quiver



bound by $\alpha_i \alpha_{i+1}$ for $i \in [p - r, p]$ and $\beta_i \beta_{i+1}$ for $i \in [q, q + r']$,

and the following series of lemmas.

Lemma 1.11. *If $p, q \in \mathbb{N}_+$, $r \in [0, p - 1]$, and $q > 1$, then $\Gamma_0(p, q, r)$ is tilting-cotilting equivalent to $\Gamma_0(p + 1, q - 1, r)$.*

Proof. It is enough to apply the generalized APR-reflection at b_{q-1} , followed by the APR-coreflection at z , the generalized APR-coreflection at y , and the APR-coreflections at b_{q-2}, \dots, b_1 (we omit the last step if $q = 2$). \square

Lemma 1.12. *If $p, q \in \mathbb{N}_+$, $r \in [0, p - 1]$, $r' \in \mathbb{N}$, and $r' \geq r$, then $\Gamma_1(p, q, r, r')$ is tilting-cotilting equivalent to $\Gamma_2(q + r' - r, p, r' - r, r)$.*


Proof. First for each $i \in [1, r]$ we apply the HW-coreflection at z followed by the APR-reflection at z , and the generalized APR-coreflection at $a_{p+r'-i}$ applied $r+r'+1-i$ times. Next we apply the HW-coreflections at $z, a_{p+r'-r-1}, \dots, a_p$ (only at z if $r = r'$) and we obtain a bound quiver whose bound quiver algebra is easily seen to be tilting-cotilting equivalent to $\Gamma_2(q + r' - r, p, r' - r, r)$. \square

Lemma 1.13. *If $p, q \in \mathbb{N}_+$, $r \in [0, p - 1]$, $r' \in \mathbb{N}$, and $r \geq r'$, then $\Gamma_1(p, q, r, r')$ is tilting-cotilting equivalent to $\Gamma_2(p + 2r' - r, q, r', r - r')$.*

Proof. Since $\Gamma_1(p, q, r, r')$ is tilting-cotilting equivalent to $\Gamma_1(p + r' - r, q, r', r)^{\text{op}}$ and $\Gamma_2(p + 2r' - r, q, r', r - r')$ is tilting-cotilting equivalent to $\Gamma_2(q + r - r', p + r' - r, r - r', r')^{\text{op}}$, hence the claim follows from the previous lemma. \square

Lemma 1.14. *If $p, q \in \mathbb{N}_+$, $r \in [0, p - 1]$, $r' \in \mathbb{N}$, and $r \geq r'$, then $\Gamma_2(p, q, r, r')$ is tilting-cotilting equivalent to $\Gamma_2(p, q, r - r', r')$.*

Proof. By applying the APR-coreflection at z followed by the generalized APR-coreflection at z applied r' times, we replace $\Gamma_2(p, q, r, r')$ by (an algebra isomorphic to) the bound quiver algebra of the quiver



bound by $\alpha'_i \alpha'_{i+1}$ for $i \in [p - r, p]$ and $\beta'_i \beta'_{i+1}$ for $i \in [q, q + r']$. It is easily seen that this algebra is tilting-cotilting equivalent to $\Gamma_2(p, q, r - r', r')$ (we just shift relations sufficiently many times). \square


Lemma 1.15. *If $p, q \in \mathbb{N}$, $r \in [0, p - 1]$, $r' \in \mathbb{N}$, and $r' \geq r$, then $\Gamma_2(p, q, r, r')$ is tilting-cotilting equivalent to $\Gamma_2(p, q + r, r, r' - r)$.*

Proof. Since $\Gamma_2(p, q, r, r')$ is tilting-cotilting equivalent to $\Gamma_2(q + r', p - r, r', r)^{\text{op}}$ and $\Gamma_2(p, q + r, r, r' - r)$ is tilting-cotilting equivalent to $\Gamma_2(q + r', p - r, r' - r, r)^{\text{op}}$, hence the claim follows from the previous lemma. \square

Proof of Proposition 1.8. Without loss of generality we may assume that $\alpha_{p_1} \alpha_{p_1+1} \in R$ and $\beta_{q_1} \beta_{q_1+1} \in R$. We first show that either $\alpha_i \alpha_{i+1} \notin R$ for all $i \in [1, p_1 - 1]$ or $\beta_i \beta_{i+1} \notin R$ for all $i \in [1, q_1 - 1]$. Assume this is not the case. In particular, $p_1, q_1 \geq 2$. By shifting relations we may assume that $\alpha_1 \alpha_2 \in R$ and $\beta_1 \beta_2 \in R$. If (Δ', R') is the quiver obtained from (Δ, R) by applying the HW-reflection at x followed by the APR-reflection at x , where $x = t\alpha_1$, then $\Delta' = \Delta$ and $R' = R \setminus \{\alpha_1 \alpha_2, \beta_1 \beta_2\}$, hence the claim follows by induction. Similarly, we prove that either $\alpha_i \alpha_{i+1} \notin R$ for all $i \in [p_1 + 1, p_1 + p_2 - 1]$ or $\beta_i \beta_{i+1} \notin R$ for all $i \in [q_1 + 1, q_1 + q_2 - 1]$. Consequently, by shifting relations one easily observes that the bound quiver algebra of (Δ, R) is tilting-cotilting equivalent either to $\Gamma_1(p, q, r, r')$ or to $\Gamma_2(p, q, r, r')$ for some $p, q \in \mathbb{N}_+$, $r \in [0, p - 1]$, and $r' \in \mathbb{N}$. Since $\Gamma_1(p, q, r, 0) = \Gamma_0(p, q, r) = \Gamma_2(p, q, r, 0)$ for all $p, q \in \mathbb{N}_+$ and $r \in \mathbb{N}$, $\Gamma_1(p, q, 0, r') \simeq \Gamma_0(p + r', q, r')^{\text{op}}$ and $\Gamma_2(p, q, 0, r') \simeq \Gamma_0(q + r', p, r')^{\text{op}}$ for all $p, q \in \mathbb{N}_+$ and $r' \in \mathbb{N}$, and $\Gamma_0(p, 1, r)$ is tilting-cotilting equivalent to $\Lambda'_0(p, r)$ for all $p \in \mathbb{N}_+$ and $r \in [0, p - 1]$, hence the claim follows from the above series of lemmas. \square

We finish this section with the following.

Proposition 1.16. *If (Δ, R) is a gentle bound quiver such that*




for some $p_1, p_2, p_3 \in \mathbb{N}_+$, and $\beta_{p_2}\alpha_1, \gamma_{p_3}\beta_1 \in R$, then the bound quiver algebra of (Δ, R) is tilting-cotilting equivalent to $\Lambda_0(p, r)$ for some $p \in \mathbb{N}_+$ and $r \in [0, p - 1]$.

Proof. Let r_1 be the number of $i \in [1, p_1 - 1]$ such that $\alpha_i\alpha_{i+1} \in R$, let r_2 be the number of $i \in [1, p_2 - 1]$ such that $\beta_i\beta_{i+1} \in R$, and let r_3 be the number of $i \in [1, p_3 - 1]$ such that $\gamma_i\gamma_{i+1} \in R$. We prove the claim by induction on $r_1 + r_2 + r_3$.


If $r_1 = 0 = r_3$, then it follows by shifting relations that the bound quiver algebra of (Δ, R) is tilting-cotilting equivalent to $\Lambda_0(p_1 + p_2 + p_3 - 2, r_2)$.

If $r_1 > 0$ and $r_3 = 0$, then by shifting relations we may assume that $p_3 = 1$ and $\alpha_1\alpha_2 \in R$. If (Δ', R') is the bound quiver obtained from (Δ, R) by applying the generalized APR-reflection at u followed by the APR-reflection at x_1 , then $R' = R \setminus \{\alpha_1\alpha_2, \beta_{p_2}\alpha_1, \gamma_1\beta_1\} \cup \{\gamma_1\alpha_2, \beta_{p_2}\gamma_1\}$ and



hence the claim follows by induction. Dually, the claim follows if $r_1 = 0$ and $r_3 > 0$.

Assume finally that $r_1 > 0$ and $r_3 > 0$. By shifting relations we may assume that $\alpha_1\alpha_2 \in R$ and $\gamma_1\gamma_2 \in R$. If (Δ', R') is obtained from (Δ, R) by applying the generalized APR-reflection at u followed by the APR-reflection at x_1 , then $R' = R \setminus \{\alpha_1\alpha_2, \beta_1\alpha_1, \gamma_1\gamma_2\} \cup \{\beta_{p_2}\gamma_1, \gamma_1\alpha_2\}$ and



and the claim again follows by induction. \square

2. COMPLETENESS OF THE LIST

We start our considerations in this section by extending the list of algebras in Theorems 1 and 2. Namely, as a consequence of the Lemmas 1.4 and 1.9 and Corollary 1.6, it follows that in order to show the completeness of the lists in Theorems 1 and 2, it is enough to prove the following.

Proposition 2.1. *If (Δ, R) is a gentle two-cycle bound quiver, then the bound quiver algebra of (Δ, R) is tilting-cotilting equivalent to one of the following algebras:*

- $\Lambda_0(p, r)$ for some $p \in \mathbb{N}_+$ and $r \in [0, p - 1]$,
- $\Lambda'_0(p, r)$ for some $p \in \mathbb{N}_+$ and $r \in [0, p - 1]$,
- $\Lambda_1(p_1, p_2, p_3, p_4, r_1)$ for some $p_1, p_2 \in \mathbb{N}_+$, $p_3, p_4 \in \mathbb{N}$, and $r_1 \in [0, p_1 - 1]$, such that $p_2 + p_3 \geq 2$ and $p_4 + r_1 \geq 1$,
- $\Lambda_2(p_1, p_2, p_3, r_1, r_2)$ for some $p_1, p_2 \in \mathbb{N}_+$, $p_3 \in \mathbb{N}$, $r_1 \in [0, p_1 - 1]$, $r_2 \in [0, p_2 - 1]$, such that $p_3 + r_1 + r_2 \geq 1$.

For the rest of the section we assume that (Δ, R) is a gentle two-cycle bound quiver. We show, in a sequence of steps, that the bound quiver algebra of (Δ, R) is tilting-cotilting equivalent to one of the algebras listed in the above proposition.

We may divide the arrows in Δ into three disjoint groups:

- $\alpha \in \Delta_1$ is called a cycle arrow if the quiver $(\Delta_0, \Delta_1 \setminus \{\alpha\})$ is connected,
- $\alpha \in \Delta_1$ is called a branch arrow if the quiver $(\Delta_0, \Delta_1 \setminus \{\alpha\})$ has a connected component which is a two-cycle quiver,
- $\alpha \in \Delta_1$ is called a connecting arrow if the quiver $(\Delta_0, \Delta_1 \setminus \{\alpha\})$ has two connected components which are one-cycle quivers.

A vertex x of Δ is called a connecting vertex if there exist at least three arrows adjacent to x which are not branch arrows. We call $\alpha\beta \in R$ a branch relation if either α or β is a branch arrow.

Step 1. *We may assume that there are no branch relations in R .*

Proof. If there exists a branch relation in (Δ, R) , then by passing, if necessary, to the opposite algebra, we may assume that there exists a subquiver

$$\Sigma = \bullet_{x_0} \xrightarrow{\alpha_1} \dots \xrightarrow{\alpha_{n-2}} \bullet_{x_{n-2}} \xleftarrow{\alpha_{n-1}} \bullet_{x_{n-1}} \xleftarrow{\alpha_n} \bullet_{x_n},$$

of Δ for some $n \geq 2$, where $\alpha_1, \dots, \alpha_{n-2}$ are free arrows, $\alpha_{n-1}\alpha_n \in R$, and there are no other arrows adjacent to x_0, \dots, x_{n-2} (in particular, α_{n-1} is a branch arrow, hence $\alpha_{n-1}\alpha_n$ is a branch relation). By applying APR-coreflections we may assume that $s\alpha_i = x_i$ for all $i \in [1, n - 2]$. If (Δ', R') is the bound quiver obtained from (Δ, R) by applying

the generalized APR-reflections at x_{n-2}, \dots, x_1 followed by the APR-reflection at x_0 , then $R' = R \setminus \{\alpha_{n-1}\alpha_n\}$ and Δ' is obtained from Δ by replacing Σ by the quiver

$$\bullet_{x_{n-1}} \xleftarrow{\alpha_1} \bullet_{x_0} \xleftarrow{\alpha_2} \dots \xleftarrow{\alpha_{n-1}} \bullet_{x_{n-2}} \xleftarrow{\alpha_n} \bullet_{x_n} .$$

In particular, the number of branch relations decreases, hence the claim follows by induction. \square

By a branch in Δ we mean a maximal nontrivial (i.e. with nonempty set of arrows) connected subquiver of Δ whose all arrows are branch arrows. We say that a branch B in Δ is rooted at x if $x \in B_0$ and there exists $\alpha \in \Delta_1$ adjacent to x which is not a branch arrow. An immediate consequence of the assumption made in the above step is that each branch B in Δ is a linear quiver rooted at one of its ends. Moreover, by applying the APR-reflections we may assume that B is equioriented and rooted at its unique sink.

Step 2. *We may assume that there are no branch arrows in Δ .*

Proof. We say that $x \in \Delta_0$ is an insertion vertex if either x is the connecting vertex or there exists $\alpha \in \Delta_1$ such that $s\alpha = x$, α is not a branch arrow, and there is no $\beta \in \Delta_1$ with $t\beta = x$ and $\alpha\beta \in R$. Observe that there is no branch which is rooted at an insertion vertex. Moreover, for each $x \in \Delta_0$ there exists a path in Δ starting at an insertion vertex and terminating at x . In particular, if B is a branch rooted at x , then we call the minimal length of such a path the distance between B and an insertion vertex. We prove our claim by induction on the number of branches in (Δ, R) and, for a given branch B , by the distance between B and an insertion vertex.

Let

$$B = \bullet_{x_0} \xleftarrow{\alpha_1} \bullet_{x_1} \xleftarrow{\dots} \bullet_{x_{n-1}} \xleftarrow{\alpha_n} \bullet_{x_n}, \quad n \in \mathbb{N}_+,$$

be a branch in Δ . Let α and β be the arrows in Δ with $s\alpha = x_0 = t\beta$ and $\beta \neq \alpha_1$. Observe that $\alpha\beta \in R$ and there are no other arrows adjacent to x_0 . Put $y = t\alpha$ and $z = s\beta$.

Assume first that there is no $\gamma \in \Delta_1$ with $t\gamma = z$ and $\beta\gamma \in R$. If (Δ', R') is the bound quiver obtained from (Δ, R) by applying the generalized APR-reflections at x_0, \dots, x_{n-1} , then $R' = R \setminus \{\alpha\beta\} \cup \{\alpha\alpha_n\}$ and Δ' is obtained from Δ by replacing the subquiver


$$\begin{array}{ccccccc} & & y & & & & \\ & & \bullet & & & & \\ & & \uparrow & & & & \\ & & \alpha & & & & \\ z & \xrightarrow{\beta} & \bullet_{x_0} & \xleftarrow{\alpha_1} & \bullet_{x_1} & \xleftarrow{\dots} & \bullet_{x_{n-1}} \xleftarrow{\alpha_n} \bullet_{x_n} \end{array}$$

by the quiver


$$z \xleftarrow{\beta} \bullet_{x_0} \xleftarrow{\alpha_1} \bullet_{x_1} \xleftarrow{\dots} \bullet_{x_{n-1}} \xrightarrow{\alpha_n} \bullet_{x_n} \xrightarrow{\alpha} \bullet_y,$$

hence the claim follows in this case.

Assume now that there exists $\gamma \in \Delta_1$ with $t\gamma = z$ and $\beta\gamma \in R$, and z is a connecting vertex in Δ_1 . Put $v = s\gamma$. If (Δ', R') is the bound quiver obtained from (Δ, R) by applying the generalized APR-reflections at x_0, \dots, x_{n-1} , then $R' = R \setminus \{\alpha\beta, \beta\gamma\} \cup \{\alpha\alpha_n, \alpha_n\gamma\}$ and Δ' is obtained from Δ by replacing the subquiver




by the quiver



Observe that the assumption that z is a connecting vertex in Δ implies that $\beta, \alpha_1, \dots, \alpha_{n-1}$ are not branch arrows in Δ' .

Finally assume that there exists $\gamma \in \Delta_1$ with $t\gamma = z$ and $\beta\gamma \in R$, but z is not a connecting vertex in Δ_1 . By induction we may assume that there is no branch rooted at z . If (Δ', R') is the bound quiver obtained from (Δ, R) by applying the HW-coreflection at x_i followed by the APR-reflection at x_i for $i = n, \dots, 1$, then $R' = R$ and Δ' is obtained from Δ by replacing the subquiver



by the quiver



and the claim follows by induction. \square

We say that Δ is special if either there is a unique connecting vertex in Δ or there is a connecting arrow in Δ . Otherwise, we call Δ proper. We concentrate now on the case when Δ is special. We describe first more precisely its structure. We may divide the cycle arrows of Δ into two disjoint subsets $\Delta_1^{(1)}$ and $\Delta_1^{(2)}$ in such a way that cycle arrows α and β belong to the same subset if and only if the quiver $(\Delta_0, \Delta_1 \setminus \{\alpha, \beta\})$ has a connected component which is a one-cycle quiver. For $j \in [1, 2]$ we denote by $\Delta^{(j)}$ the minimal subquiver of Δ with the set of arrows $\Delta_1^{(j)}$. Observe that $\Delta^{(j)}$ is a (not necessarily oriented) cycle. We divide the arrows in $\Delta^{(j)}$ into disjoint subsets $\Delta_{1,-}^{(j)}$ and $\Delta_{1,+}^{(j)}$ in such a way that if α and β are adjacent to the same vertex for $\alpha, \beta \in \Delta_1^{(j)}$, $\alpha \neq \beta$,

then α and β belong to the same subset if and only if either $s\alpha = t\beta$ or $t\alpha = s\beta$. For $\varepsilon \in \{-, +\}$ we put

$$R_\varepsilon^{(j)} = \{\alpha\beta \in R \mid \alpha, \beta \in \Delta_{1,\varepsilon}^{(j)}\}.$$

Step 3. *If Δ is special, then we may assume that for each $j \in [1, 2]$ there exists $\varepsilon \in \{-, +\}$ such that $R_\varepsilon^{(j)} = \emptyset$.*

Proof. If $\Delta^{(j)}$ is an oriented cycle, then there is nothing to prove, hence assume that $\Delta^{(j)}$ is not an oriented cycle and $R_-^{(j)} \neq \emptyset \neq R_+^{(j)}$. There exists a subquiver

$$\Sigma = \bullet_{y_1} \xleftarrow{\alpha_1} \bullet_{y_2} \xleftarrow{\alpha_2} \bullet_{x_0} \xrightarrow{\gamma_1} \dots \xrightarrow{\gamma_n} \bullet_{x_n} \xrightarrow{\beta_2} \bullet_{z_2} \xrightarrow{\beta_1} \bullet_{z_1}$$

of Δ for some $n \in \mathbb{N}$, such that $\alpha_1\alpha_2 \in R_-^{(j)}$, $\beta_1\beta_2 \in R_+^{(j)}$, there are no other arrows adjacent to x_0, \dots, x_n , and $\gamma_1, \dots, \gamma_n$ are free arrows. By applying appropriate APR-reflections at x_1, \dots, x_{n-1} (see the discussion after Lemma 1.3) we may assume that

$$\Sigma = \bullet_{y_1} \xleftarrow{\alpha_1} \bullet_{y_2} \xleftarrow{\alpha_2} \bullet_{x_0} \xleftarrow{\gamma_1} \dots \xleftarrow{\gamma_k} \bullet_{x_k} \xrightarrow{\gamma_{k+1}} \dots \xrightarrow{\gamma_n} \bullet_{x_n} \xrightarrow{\beta_2} \bullet_{z_2} \xrightarrow{\beta_1} \bullet_{z_1}$$

for some $k \in [0, n]$. By shifting the relations $\alpha_1\alpha_2$ and $\beta_1\beta_2$ to the right, we may assume that $n = 0$, i.e.

$$\Sigma = \bullet_{y_1} \xleftarrow{\alpha_1} \bullet_{y_2} \xleftarrow{\alpha_2} \bullet_x \xrightarrow{\beta_2} \bullet_{z_2} \xrightarrow{\beta_1} \bullet_{z_1}.$$

Assume first that neither y_2 nor z_2 is a connecting vertex. If (Δ', R') is the bound quiver obtained from (Δ, R) by applying the APR-coreflections at x, y_2 , and z_2 , then $R' = R \setminus \{\alpha_1\alpha_2, \beta_1\beta_2\}$ and Δ' is obtained from Δ by replacing Σ by the quiver

$$\bullet_{y_1} \xleftarrow{\alpha_1} \bullet_{z_2} \xleftarrow{\beta_2} \bullet_x \xrightarrow{\alpha_2} \bullet_{y_2} \xrightarrow{\beta_1} \bullet_{z_1},$$

and the claim follows by induction. Otherwise, we may assume without loss of generality that y_2 is a connecting vertex and z_2 is not a connecting vertex. If (Δ', R') is the bound quiver obtained from (Δ, R) by applying the APR-coreflections at x and z_2 , then $R' = R \setminus \{\alpha_1\alpha_2, \beta_1\beta_2\} \cup \{\beta_1\alpha_2\}$ and Δ' is obtained from Δ by replacing Σ by the quiver

$$\begin{array}{ccccc} & & y_2 & & \\ & & \bullet & & \\ & & \downarrow \alpha_2 & & \\ \bullet_{y_1} & \xleftarrow{\alpha_1} & \bullet_{z_2} & \xleftarrow{\beta_2} & \bullet_x & \xrightarrow{\beta_1} & \bullet_{z_1} \end{array}.$$

Observe that α_2 is a connecting arrow in Δ' , hence the claim again follows by induction. \square


Step 4. *If Δ is special, then for each $j \in [1, 2]$ we may assume that $\Delta^{(j)}$ is either an oriented cycle or there is a unique source (equivalently, unique sink) in $\Delta^{(j)}$.*

Proof. This follows easily by applying APR-reflections and shifts of relations (see the discussion after Lemma 1.3). \square


Step 5. *If Δ is special, then we may assume that either there is no connecting arrow in Δ or, for each $j \in [1, 2]$, $\Delta^{(j)}$ is an oriented cycle and $\alpha\beta \in R$ for all $\alpha, \beta \in \Delta_1^{(j)}$ with $s\alpha = t\beta$.*

Proof. We prove the claim by induction on the sum of the number of connecting arrows and the number of connecting relations, where we say that $\alpha\beta \in R$ is a connecting relation if both α and β are connecting arrows. We may assume without loss of generality that either $\Delta^{(1)}$ is not an oriented cycle or there exist $\alpha, \beta \in \Delta_1^{(1)}$ with $s\alpha = t\beta$ and $\alpha\beta \notin R$. Let $x \in \Delta_0^{(1)}$ be a connecting vertex. Let α be the connecting arrow adjacent to x . Without loss of generality we may assume that $x = s\alpha$. Let β and γ be the arrows adjacent to x different from α . Again we may assume without loss of generality that $x = t\beta$. By symmetry we may also assume that $\alpha\beta \in R$ if $x = t\gamma$. Put $y = t\alpha$ and $z = s\beta$. In order to make it easier to track the proof we will number the cases.

(1) Assume that $\alpha\beta \notin R$. According to our assumptions this implies that $x = s\gamma$ and $\gamma\beta \in R$. Put $v = t\gamma$. If $\Delta^{(1)}$ is not an oriented cycle, then by applying APR-reflections and the dual of Lemma 1.2 we may assume that v is a sink. In particular, there is no $\gamma' \in \Delta_1$ with $s\gamma' = v$ and $\gamma'\gamma \in R$. By shifting relations we may also assume that this condition is satisfied if $\Delta^{(1)}$ is an oriented cycle. Let (Δ', R') be the bound quiver obtained from (Δ, R) by applying the generalized APR-coreflection at x . If there is no $\alpha' \in \Delta_1$ with $s\alpha' = y$ and $\alpha'\alpha \in R$, then $R' = R \setminus \{\gamma\beta\} \cup \{\alpha\beta\}$ and Δ' is obtained from Δ by replacing the subquiver




by the quiver




On the other hand, if there exists $\alpha' \in \Delta_1$ with $s\alpha' = y$ and $\alpha'\alpha \in R$, then $R = R \setminus \{\gamma\beta, \alpha'\alpha\} \cup \{\alpha\beta, \alpha'\gamma\}$ and Δ' is obtained from Δ by

replacing the subquiver



by the quiver




where $y' = t\alpha'$. Observe, that either Δ' is proper (if y is a connecting vertex in the second case) or we decrease the number of connecting arrows (otherwise), hence the claim follows by induction.

(2) Assume that $\alpha\beta \in R$.


(2.1) Assume that there is no $\alpha' \in \Delta'_1$ with $s\alpha' = y$ and $\alpha'\alpha \in R$.

(2.1.1) Assume that y is a connecting vertex. If either $\Delta^{(2)}$ is not an oriented cycle or there exist $\delta', \delta'' \in \Delta_1^{(2)}$ with $s\delta' = t\delta''$ and $\delta'\delta'' \notin R$, then the claim follows by symmetry from (1), thus we may assume that $\Delta^{(2)}$ is an oriented cycle such that $\delta'\delta'' \in R$ for all $\delta', \delta'' \in \Delta_1^{(2)}$ with $s\delta' = t\delta''$.

(2.1.1.1) Assume that $|\Delta_1^{(2)}| = 1$. If (Δ', R') is the bound quiver obtained from (Δ, R) by applying the generalized APR-reflection at y , then $R' = R$ and Δ' is obtained from Δ by replacing the subquiver




by the quiver



hence the claim follows.

(2.1.1.2) Assume that $|\Delta_1^{(2)}| > 1$. Let α' and β' be the arrows in $\Delta^{(2)}$ with $s\alpha' = y = t\beta'$. Put $v' = t\alpha'$ and $x' = s\beta'$. Let γ' be the arrow in $\Delta^{(2)}$ with $t\gamma' = x'$. Put $z' = s\gamma'$. Recall that $\alpha'\beta', \beta'\gamma' \in R$. If (Δ', R') is the bound quiver obtained from (Δ, R) by applying the generalized APR-reflection at y followed by the APR-reflection at x' , then $R' = R \setminus \{\alpha\beta, \alpha'\beta', \beta'\gamma'\} \cup \{\alpha\gamma', \alpha'\alpha\}$ and Δ' is obtained from Δ by replacing the subquiver



by the quiver

$$\begin{array}{ccccccc} & & & & z' & & \\ & & & & \downarrow \gamma' & & \\ z & \xrightarrow{\beta} & x' & \xrightarrow{\beta'} & y & \xrightarrow{\alpha} & x & \xrightarrow{\alpha'} & v' \end{array},$$

hence the claim follows in this case.

(2.1.2) Assume that y is not a connecting vertex.

(2.1.2.1) Assume that there exists $\alpha' \in \Delta_1$ with $s\alpha' = y$. Our assumptions imply that $\alpha'\alpha \notin R$. Put $y' = t\alpha'$. If (Δ', R') is the bound quiver obtained from (Δ, R) by applying the generalized APR-reflection at y , then $R' = R \setminus \{\alpha\beta\} \cup \{\alpha'\alpha\}$ and Δ' is obtained from Δ by replacing the subquiver

$$z \xrightarrow{\beta} x \xrightarrow{\alpha} y \xrightarrow{\alpha'} y'$$

by the quiver

$$z \xrightarrow{\beta} y \xrightarrow{\alpha} x \xrightarrow{\alpha'} y',$$

hence the claim follows by induction.

(2.1.2.2) Assume there exists $\alpha' \in \Delta'$ with $t\alpha' = y$. Put $x' = s\alpha'$.

(2.1.2.2.1) Assume that either x' is a connecting vertex or α' is a free arrow. Moreover, if x' is a connecting arrow and α' is not a free arrow, then let β' be the arrow in Δ with $t\beta' = x'$ and $\alpha'\beta' \in R$, and put $z' = s\beta'$. Let (Δ', R') be the bound quiver obtained from (Δ, R) by applying the APR-reflection at y . If α' is a free arrow, then $R' = R \setminus \{\alpha\beta\} \cup \{\alpha'\beta\}$ and Δ' is obtained from Δ by replacing the subquiver

$$z \xrightarrow{\beta} x \xrightarrow{\alpha} y \xleftarrow{\alpha'} x'$$

by the quiver

$$\begin{array}{ccc} & z & \\ & \downarrow \beta & \\ x & \xleftarrow{\alpha} & y & \xrightarrow{\alpha'} & x' \end{array},$$

hence the claim follows by induction. On the other hand, if x' is a connecting arrow and α' is not a free arrow, then $R' = R \setminus \{\alpha\beta, \alpha'\beta'\} \cup \{\alpha\beta', \alpha'\beta\}$ and Δ' is obtained from Δ by replacing the subquiver

$$z \xrightarrow{\beta} x \xrightarrow{\alpha} y \xleftarrow{\alpha'} x' \xleftarrow{\beta'} z'$$

by the quiver

$$\begin{array}{ccc} z & & z' \\ & \searrow \beta & \swarrow \beta' \\ x & \xleftarrow{\alpha} & y & \xrightarrow{\alpha'} & x' \end{array},$$

hence the claim follows.

(2.1.2.2.2) Assume that x' is not a connecting vertex and there exists $\beta' \in \Delta_1$ with $t\beta' = x'$ and $\alpha'\beta' \in R$. Put $z' = s\beta'$. If (Δ', R') is the bound quiver obtained from (Δ, R) by applying the APR-reflections at y and x' , then $R' = R \setminus \{\alpha\beta, \alpha'\beta'\} \cup \{\alpha\beta'\}$ and Δ' is obtained from Δ by replacing the subquiver

$$\bullet \xrightarrow{\beta} \bullet \xrightarrow{\alpha} \bullet \xleftarrow{\alpha'} \bullet \xleftarrow{\beta'} \bullet$$

$z \quad x \quad y \quad x' \quad z'$

by the quiver

$$\begin{array}{ccccc} & & x & & \\ & & \bullet & & \\ & & \uparrow \alpha & & \\ \bullet & \xrightarrow{\beta} & \bullet & \xleftarrow{\alpha'} & \bullet \\ z & & x' & & y & & z' \end{array},$$

hence the claim follows by induction.

(2.2) Assume that there exists $\alpha' \in \Delta_1$ with $s\alpha' = y$ and $\alpha'\alpha \in R$. Put $y' = t\alpha'$.

(2.2.1) Assume that $x = t\gamma$. Let $\beta_1 \cdots \beta_n$ and $\gamma_1 \cdots \gamma_m$ be the maximal paths in Δ terminating at x with $\beta_1 = \beta$ and $\gamma_1 = \gamma$. Put $u = s\beta_n$, $u'_i = s\beta_i$ for $i \in [1, n-1]$ and $u''_i = s\gamma_i$ for $i \in [1, m-1]$.

(2.2.1.1) Assume that there exists $i \in [1, m-1]$ such that $\gamma_i\gamma_{i+1} \in R$. By shifting relations we may assume that $\gamma_{m-1}\gamma_m \in R$. Observe that $\beta_i\beta_{i+1} \notin R$ for all $i \in [1, n-1]$. If (Δ', R') is the bound quiver obtained from (Δ, R) by applying the HW-coreflection at u followed by the composition of the HW-coreflection at u'_i and the APR-reflection at u'_i for $i = n-1, \dots, 1$, then $R' = R \setminus \{\gamma_{m-1}\gamma_m, \alpha\beta\} \cup \{\beta_n\gamma\}$ and Δ' is obtained from Δ by replacing the subquiver

$$\bullet \xleftarrow{\gamma_m} \bullet \xrightarrow{\beta_n} \bullet \cdots \xrightarrow{\beta_1} \bullet$$

$u''_{m-1} \quad u \quad u'_{n-1} \quad \dots \quad u'_1 \quad x$

by the quiver

$$\bullet \xrightarrow{\gamma_m} \bullet \cdots \xrightarrow{\beta_1} \bullet \xrightarrow{\beta_n} \bullet \xleftarrow{\beta_n} \bullet$$

$u''_{m-1} \quad u'_{n-1} \quad \dots \quad u'_1 \quad u \quad x$

hence we reduce the proof to (1).

(2.2.1.2) Assume that $\gamma_i\gamma_{i+1} \notin R$ for all $i \in [1, m-1]$. Let r be the number of $i \in [1, n-1]$ such that $\beta_i\beta_{i+1} \in R$. By shifting relations we may assume $\beta_i\beta_{i+1} \in R$ for all $i \in [n-r, n-1]$. Put $\beta_0 = \alpha$. If (Δ', R') is the bound quiver obtained from (Δ, R) by applying the generalized APR-coreflections at u'_1, \dots, u'_{n-r-1} , then $R' = R \setminus \{\alpha\beta\} \cup \{\beta_{n-r-1}\beta_{n-r}\}$ and Δ' is obtained from Δ by replacing the subquiver

$$\bullet \xrightarrow{\beta_n} \cdots \xrightarrow{\beta_1} \bullet \xrightarrow{\beta_0} \bullet$$

$u \quad \dots \quad x \quad y$

by the quiver

$$\bullet \xrightarrow{\beta_n} \dots \xrightarrow{\beta_{n-r}} \bullet \xrightarrow{\beta_{n-r-1}} \dots \xrightarrow{\beta_0} \bullet \xrightarrow{\beta_0} \bullet .$$

Let $\gamma'_1 \cdots \gamma'_l$ be the maximal path in (Δ', R') with $\gamma'_l = \alpha$. Observe that $l > 1$ implies that y is a connecting vertex. Put

$$u' = \begin{cases} u'_{n-1} & r \geq 1, \\ x & r = 0 \end{cases} \quad \text{and} \quad v' = t\gamma'_1.$$

Let (Δ'', R'') be the bound quiver obtained from (Δ', R') by applying the HW-coreflection at u followed by the composition of the HW-coreflection at u''_i and the APR-reflection at u''_i for $i = m-1, \dots, 1$. If there exists δ in Δ with $t\delta = v'$ and $\delta \neq \gamma'_1$, then $R'' = R' \setminus \{\beta_{n-1}\beta_n\} \cup \{\gamma_m\delta\}$, while $R'' = R' \setminus \{\beta_{n-1}\beta_n\}$, otherwise. Moreover, Δ'' is obtained from Δ' by replacing the subquiver

$$\bullet \xleftarrow{\beta_n} \bullet \xrightarrow{\gamma_m} \bullet \xrightarrow{\gamma_{m-1}} \dots \xrightarrow{\gamma_1} \bullet \xrightarrow{\beta_{n-r-1}} \dots \xrightarrow{\beta_0} \bullet \xrightarrow{\gamma'_{l-1}} \dots \xrightarrow{\gamma'_1} \bullet$$

by the quiver

$$\bullet \xrightarrow{\beta_n} \bullet \xrightarrow{\gamma_{m-1}} \dots \xrightarrow{\gamma_1} \bullet \xleftarrow{\gamma_m} \bullet \xleftarrow{\gamma'_1} \dots \xleftarrow{\gamma'_{l-1}} \bullet \xleftarrow{\beta_0} \dots \xleftarrow{\beta_{n-r-1}} \bullet ,$$

and the claim follows (by induction if y is not a connecting vertex).

(2.2.2) Assume that $x = s\gamma$. Put $v = t\gamma$.

(2.2.2.1) Assume that there exists $\gamma' \in \Delta_1$ with $s\gamma' = v$ and $\gamma'\gamma \in R$ (by shifting relations we may assume that this condition is satisfied if $\Delta^{(1)}$ is an oriented cycle). Put $v' = t\gamma'$. If (Δ', R') is the bound quiver obtained from (Δ, R) by applying the generalized coreflection at x followed, if y is not a connecting vertex, by the APR-coreflection at y , then

$$R' = \begin{cases} R \setminus \{\alpha'\alpha, \alpha\beta, \gamma'\gamma\} \cup \{\gamma'\alpha, \gamma\beta, \alpha'\gamma\} & y \text{ is a connecting vertex,} \\ R \setminus \{\alpha'\alpha, \alpha\beta, \gamma'\gamma\} \cup \{\gamma\beta, \alpha'\gamma\} & y \text{ is not a connecting vertex,} \end{cases}$$


and Δ' is obtained from Δ by replacing the subquiver

$$\begin{array}{ccccccc} & & z & & & & \\ & & \bullet & & & & \\ & & \downarrow \beta & & & & \\ v' & \xleftarrow{\gamma'} & v & \xleftarrow{\gamma} & x & \xrightarrow{\alpha} & y & \xrightarrow{\alpha'} & y' \end{array}$$

by the quiver

$$\begin{array}{ccccccc} & & v' & & & & \bullet \\ & & \downarrow \gamma' & & \uparrow \alpha' & & \\ z & \xrightarrow{\beta} & v & \xrightarrow{\gamma} & x & \xleftarrow{\alpha} & y \\ & & & & & & \bullet \end{array}$$

if y is a connecting vertex, or



if y is not a connecting vertex, hence the claim again follows.


(2.2.2.2) Assume that $\Delta^{(1)}$ is not an oriented cycle. Let $\gamma_1 \cdots \gamma_n$ be the maximal path in Δ with $\gamma_n = \gamma$. We may additionally assume that $\gamma_i \gamma_{i+1} \notin R$ for all $i \in [1, n-1]$. Consequently, we may reduce the proof in this case to (2.2.1) by applying APR-reflections and shifts of relations. \square

Step 6. *If Δ is special, then we may assume that for each $j \in [1, 2]$ $\Delta^{(j)}$ is an oriented cycle or either the source or the sink in $\Delta^{(j)}$ is a connecting vertex.*

Proof. If both $\Delta^{(1)}$ and $\Delta^{(2)}$ are not oriented cycles, then there is nothing to prove, thus without loss of generality we may assume that $\Delta^{(1)}$ is not an oriented cycle. Observe that our assumptions imply that there are no connecting arrows in Δ . Let x be the connecting vertex in Δ and assume that x is neither a source nor a sink in $\Delta^{(1)}$. Observe that $x \in \Delta_0^{(1)} \cap \Delta_0^{(2)}$. Let α, β, α' and β' be the arrows in Δ with $s\alpha = t\beta = x = s\alpha' = t\beta'$, $\alpha, \beta \in \Delta_1^{(1)}$, and $\alpha', \beta' \in \Delta_1^{(2)}$. Put $y = t\alpha$, $y' = t\alpha'$, $z = s\beta$, and $z' = s\beta'$. By applying APR-coreflections, shifts of relations and Lemma 1.2 we may assume that z is a source in $\Delta^{(1)}$.

Assume first that $\alpha' = \beta'$. Then $\alpha\beta \in R$ and $\alpha'\beta' \in R$. Let $\gamma_1 \cdots \gamma_m$ be the maximal path in Δ starting at z with $\gamma_m \neq \beta$. Observe that $\gamma_i \gamma_{i+1} \notin R$ for all $i \in [1, m-1]$. Put $v_i = s\gamma_i$ for $i \in [1, m-1]$. The bound quiver algebra of the bound quiver obtained from (Δ, R) by applying the APR-coreflections at z, v_{m-1}, \dots, v_1 , is easily seen to be tilting-cotilting equivalent to $\Lambda_2(p, 1, m, r, 0)$ for some $p \in \mathbb{N}_+$ and $r \in [0, p-1]$, hence the claim follows in this case.

Assume now that $\alpha\beta \in R$ and $\alpha'\beta' \in R$, but $\alpha' \neq \beta'$. Let (Δ', R') be the bound quiver obtained from (Δ, R) by applying the generalized APR-reflection at x . If there exists $\beta'' \in \Delta_1$ with $t\beta'' = z'$ and $\beta'\beta'' \in R$, then $R' = R \setminus \{\alpha\beta, \alpha'\beta', \beta'\beta''\} \cup \{\alpha\beta', \alpha'\beta, \beta\beta''\}$ and Δ' is obtained from Δ by replacing the subquiver



by the quiver

$$\begin{array}{ccccccc}
 & & z'' & & & & \\
 & & \downarrow \beta'' & & & & \\
 & & x & & & & \\
 \bullet & \xleftarrow{\alpha} & \bullet & \xleftarrow{\beta'} & \bullet & \xrightarrow{\beta} & \bullet & \xrightarrow{\alpha'} & \bullet \\
 y & & z' & & x & & z & & y'
 \end{array}$$

where $z'' = s\beta''$. Otherwise, $R' = R \setminus \{\alpha\beta, \alpha'\beta'\} \cup \{\alpha\beta', \alpha'\beta\}$ and Δ' is obtained from Δ by replacing the subquiver

$$\begin{array}{ccc}
 y & & y' \\
 & \swarrow \alpha & \nearrow \alpha' \\
 z & \xrightarrow{\beta} & x & \xleftarrow{\beta'} & z'
 \end{array}$$

by the quiver

$$\bullet \xleftarrow{\alpha} \bullet \xleftarrow{\beta'} \bullet \xrightarrow{\beta} \bullet \xrightarrow{\alpha'} \bullet$$

In particular, in both cases Δ' proper.

Assume finally that $\alpha\beta' \in R$ and $\alpha'\beta \in R$. Let γ be the arrow in Δ with $s\gamma = z$ and $\gamma \neq \beta$. Put $v = t\gamma$. If there exists $\gamma' \in \Delta_1$ with $s\gamma' = v$ and $\gamma'\gamma \in R$, then let (Δ', R') be the bound quiver obtained from (Δ, R) by applying the APR-coreflections at z and v , and let $v' = t\gamma'$. Observe that $R' = R \setminus \{\alpha'\beta, \gamma'\gamma\} \cup \{\gamma'\beta\}$ and Δ' is obtained from Δ by replacing the subquiver

$$\bullet \xleftarrow{\gamma'} \bullet \xleftarrow{\gamma} \bullet \xrightarrow{\beta} \bullet \xrightarrow{\alpha'} \bullet$$

by the quiver

$$\begin{array}{ccccccc}
 & & x & & & & \\
 & & \downarrow \beta & & & & \\
 & & z & & & & \\
 \bullet & \xleftarrow{\gamma'} & \bullet & \xrightarrow{\gamma} & \bullet & \xrightarrow{\alpha'} & \bullet \\
 v' & & z & & v & & y'
 \end{array}$$

Otherwise, if (Δ', R') is the bound quiver obtained from (Δ, R) by applying the APR-coreflection at z , then $R' = R \setminus \{\alpha'\beta\} \cup \{\alpha'\gamma\}$ and Δ' is obtained from Δ by replacing the subquiver

$$\bullet \xleftarrow{\gamma} \bullet \xrightarrow{\beta} \bullet \xrightarrow{\alpha'} \bullet$$

by the quiver

$$\begin{array}{ccc}
 & & y' \\
 & & \uparrow \alpha' \\
 \bullet & \xrightarrow{\gamma} & \bullet & \xleftarrow{\beta} & \bullet \\
 v & & z & & x
 \end{array}$$

Again in both cases Δ' is proper and this finishes the proof. \square

Step 7. We may assume that (Δ, R) is proper.

Proof. If $\Delta^{(1)}$ is not an oriented cycle, then also $\Delta^{(2)}$ is not an oriented cycle and the claim follows from Proposition 1.8, thus assume that $\Delta^{(1)}$ (and consequently also $\Delta^{(2)}$) is an oriented cycle.

Assume first that there are no connecting arrows in Δ and let x be the connecting vertex in Δ . Let α, β, α' and β' be the arrows in Δ with $s\alpha = t\beta = x = s\alpha' = t\beta'$, $\alpha, \beta \in \Delta_1^{(1)}$, and $\alpha', \beta' \in \Delta_1^{(2)}$. If $\alpha\beta \in R$ and $\alpha'\beta' \in R$, then it follows by shifting relations that the bound quiver algebra of (Δ, R) is tilting-cotilting equivalent to $\Lambda_2(p_1, p_2, 0, r_1, r_2)$ for some $p_1, p_2 \in \mathbb{N}_+$, $r_1 \in [0, p_1 - 1]$, and $r_2 \in [0, p_2 - 1]$, such that $r_1 + r_2 \geq 1$. On the other hand, if $\alpha\beta' \in R$ and $\alpha'\beta \in R$, then it follows by shifting relations that the bound quiver algebra of (Δ, R) is tilting-cotilting equivalent to $\Lambda_1(p_1, p_2, 0, 0, r_1)$ for some $p_1, p_2 \in \mathbb{N}_+$, $p_1, p_2 \geq 2$, and $r_1 \in [1, p_1 - 1]$.

Now assume that there are connecting arrows in Δ . Recall that in this case $\alpha\beta \in R$ for all cycle arrows α and β with $s\alpha = t\beta$. Let $\Delta^{(0)}$ be the minimal subquiver of Δ with the set of arrows consisting of the connecting arrows. Let $x \in \Delta_0^{(1)}$ and $y \in \Delta_0^{(2)}$ be the connecting vertices. Observe that $\Delta^{(0)}$ is a linear quiver. We show that we may assume that x is a unique sink in $\Delta^{(0)}$, y is a unique source in $\Delta^{(0)}$, and there are no $\alpha, \beta \in \Delta_1^{(0)}$ with $s\alpha = t\beta$ and $\alpha\beta \in R$. This will immediately imply that the bound quiver algebra of (Δ, R) is tilting-cotilting equivalent to $\Lambda_2(p_1, p_2, p_3, p_1 - 1, p_2 - 1)$ for some $p_1, p_2, p_3 \in \mathbb{N}_+$.


By repeating arguments from the proofs of Steps 3 and 4 and passing, if necessary, to the opposite algebra, we may assume that

$$\Delta = \bullet_x \xleftarrow{\alpha_1} \bullet_{x_1} \xleftarrow{\dots} \bullet_{x_{n-1}} \xleftarrow{\alpha_n} \bullet_z \xrightarrow{\beta_m} \bullet_{y_{m-1}} \xrightarrow{\dots} \bullet_{y_1} \xrightarrow{\beta_1} \bullet_y$$


for some $n \in \mathbb{N}_+$ and $m \in \mathbb{N}$, and $\beta_i\beta_{i+1} \notin R$ for all $i \in [1, m - 1]$. It is enough to show that we may additionally assume that $\alpha_i\alpha_{i+1} \notin R$ for all $i \in [1, n - 1]$, since then the claim follows from Lemma 1.5. Assume this is not the case. By shifting relations we may assume that $\alpha_1\alpha_2 \in R$.

If $|\Delta_1^{(1)}| = 1$ and (Δ', R') is the bound quiver obtained from (Δ, R) by applying the generalized APR-reflection at x followed by the APR-reflection at x_1 , then $R' = R \setminus \{\alpha_1\alpha_2\}$ and $\Delta' = \Delta$. Otherwise, let γ, δ and δ' be the arrows in $\Delta^{(1)}$ with $s\gamma = x = t\delta$ and $t\delta' = s\delta$. Observe that our assumptions imply that $\gamma\delta, \delta\delta' \in R$. Put $u = t\gamma$, $v = s\delta$ and $v' = s\delta'$. If (Δ', R') is the bound quiver obtained from (Δ, R) by applying the generalized APR-reflection at x followed by the APR-reflection at v , then $R' = R \setminus \{\gamma\delta, \delta\delta', \alpha_1\alpha_2\} \cup \{\gamma\alpha_1, \alpha_1\delta'\}$ and Δ'

is obtained from Δ by replacing the subquiver



by the quiver



where $x_2 = z$ if $n = 2$. Consequently, in both cases the claim follows by induction. \square

We investigate now the case when Δ is proper. In this case we may divide the arrows in Δ into three disjoint subsets $\Delta_1^{(1)}, \Delta_1^{(2)}, \Delta_1^{(3)}$ in such a way that $\alpha, \beta \in \Delta_1$ belong to the same subset if and only if the quiver $(\Delta_0, \Delta_1 \setminus \{\alpha, \beta\})$ has a connected component which is a one-cycle quiver. For $j \in [1, 3]$ we denote by $\Delta^{(j)}$ the minimal subquiver of Δ with the set of arrows $\Delta_1^{(j)}$. Observe that $\Delta^{(j)}$ is a linear quiver. We divide the arrows in $\Delta^{(j)}$ into disjoint subsets $\Delta_{1,-}^{(j)}$ and $\Delta_{1,+}^{(j)}$ in such a way that if α and β are adjacent to the same vertex for $\alpha, \beta \in \Delta_1^{(j)}$, $\alpha \neq \beta$, then α and β belong to the same subset if and only if either $s\alpha = t\beta$ or $t\alpha = s\beta$. For $j \in [1, 3]$ and $\varepsilon \in \{-, +\}$ we put

$$R_\varepsilon^{(j)} = \{\alpha\beta \in R \mid \alpha, \beta \in \Delta_{1,\varepsilon}^{(j)}\}.$$

Step 8. *We may assume that either $R_+^{(j)} = \emptyset$ or $R_-^{(j)} = \emptyset$ for each $j \in [1, 3]$.*

Proof. Analogous to the proof of Step 3. \square

Step 9. *We may assume that either there is at most one sink in $\Delta^{(j)}$ or there is at most one source in $\Delta^{(j)}$ for each $j \in [1, 3]$*

Proof. We prove the claim by induction on $|R|$ and, for a fixed j , on $|\Delta_1^{(j)}|$. Fix $j \in [1, 3]$ and assume that there is either a unique source or a unique sink in $\Delta^{(l)}$ for each $l \in [1, j-1]$. Let u and v be the connecting vertices in Δ , and let α and β be the arrows in $\Delta^{(j)}$ adjacent to u and v , respectively. The claim follows by the arguments presented after Lemma 1.3, unless the following condition (or its dual) is satisfied: $s\alpha = u$, $t\beta = v$, there exists $\alpha' \in \Delta_1$ with $t\alpha' = u$ and $\alpha\alpha' \in R$, and there exists $\beta' \in \Delta_1$ with $s\beta' = v$ and $\beta'\beta \in R$. Assume the above condition is satisfied. Put $x = s\beta$ and $v' = t\beta'$. If $\Delta^{(j)}$ is not an equioriented linear quiver, then by applying APR-coreflections, shifts of relations, and Lemma 1.2, we may assume that there exists $\gamma \in \Delta_1$ with $\gamma \neq \beta$ and $s\gamma = x$. Put $y = t\gamma$.

Assume there exists $\gamma' \in \Delta_1$ with $s\gamma' = y$ and $\gamma'\gamma \in R$. Put $y' = t\gamma'$. If (Δ', R') is the bound quiver obtained from (Δ, R) by applying the APR-coreflections at x and y , then $R' = R \setminus \{\beta'\beta, \gamma'\gamma\} \cup \{\gamma'\beta\}$ and Δ' is obtained from Δ by replacing the subquiver

$$\bullet_{y'} \xleftarrow{\gamma'} \bullet_y \xleftarrow{\gamma} \bullet_x \xrightarrow{\beta} \bullet_v \xrightarrow{\beta'} \bullet_{v'}$$

by the quiver

$$\begin{array}{ccccc} & & v & & \\ & & \bullet & & \\ & & \downarrow \beta & & \\ \bullet_{y'} & \xleftarrow{\gamma'} & \bullet_x & \xrightarrow{\gamma} & \bullet_y \xrightarrow{\beta'} \bullet_{v'} \end{array}$$

In particular, $|R'| < |R|$, hence the claim follows by induction in this case.

Otherwise, if (Δ', R') is the bound quiver obtained from (Δ, R) by applying the APR-coreflection at x , then $R' = R \setminus \{\beta'\beta\} \cup \{\beta'\gamma\}$ and Δ' is obtained from Δ by replacing the subquiver

$$\bullet_y \xleftarrow{\gamma} \bullet_x \xrightarrow{\beta} \bullet_v \xrightarrow{\beta'} \bullet_{v'}$$

by the quiver


$$\begin{array}{ccccc} & & v & & \\ & & \bullet & & \\ & & \downarrow \beta & & \\ \bullet_y & \xrightarrow{\gamma} & \bullet_x & \xrightarrow{\beta'} & \bullet_{v'} \end{array}$$

Observe that if $l \in [1, j-1]$ and there is no $\delta \in \Delta_1^{(l)}$ with $s\delta = v$ and $\delta \neq \beta'$, then there is either a unique source or a unique sink in $\Delta'^{(l)}$. On the other hand, if there exists such an arrow, then we may assume that there is either a unique source or a unique sink in $\Delta'^{(l)}$, since β is a free arrow in (Δ', R') . In particular, in both cases the claim follows again by induction, since $|\Delta_1^{(j)}| < |\Delta_1^{(j)}|$. \square


Step 10. *We may assume that if either $s\alpha = x = s\beta$ or $t\alpha = x = t\beta$ for a connecting vertex x , $\alpha \in \Delta_{1,\varepsilon_1}^{(j_1)}$, and $\beta \in \Delta_{1,\varepsilon_2}^{(j_2)}$, with $j_1 \neq j_2$ and $\varepsilon_1, \varepsilon_2 \in \{-, +\}$, then either $R_{\varepsilon_1}^{(j_1)} = \emptyset$ or $R_{\varepsilon_2}^{(j_2)} = \emptyset$.*

Proof. Without loss of generality we may assume that $s\alpha = x = s\beta$. If $R_{\varepsilon_1}^{(j_1)} \neq \emptyset$ or $R_{\varepsilon_2}^{(j_2)} \neq \emptyset$, then by shifting relations we may assume that there exist arrows α' and β' in Δ with $s\alpha' = t\alpha$, $s\beta' = t\beta$, and $\alpha'\alpha, \beta'\beta \in R$. Let γ be the arrow in Δ with $t\gamma = x$. Without loss of generality we may assume that $\alpha\gamma \in R$ and $\beta\gamma \notin R$. Put $y' = t\alpha'$, $z' = t\beta'$, and $u = s\gamma$. If (Δ', R') is the bound quiver obtained from (Δ, R) by applying the generalized APR-coreflections at x and y , then

$R' = R \setminus \{\alpha'\alpha, \beta'\beta, \alpha\gamma\} \cup \{\alpha'\beta, \beta\gamma\}$ and Δ' is obtained from Δ by replacing the subquiver



by the quiver



In particular, $|R'_{\varepsilon_1}(j_1)| < |R_{\varepsilon_1}(j_1)|$ and $|R'_{\varepsilon_2}(j_2)| < |R_{\varepsilon_2}(j_2)|$, hence the claim follows by induction. \square

Step 11. *We may assume that there exists $j \in [1, 3]$ such that $\Delta^{(j)}$ is equioriented.*

Proof. If the above condition is not satisfied, then without loss of generality we may assume that

$$\begin{aligned}
 \Delta^{(1)} &= \bullet \xleftarrow{\alpha_1} x_1 \xleftarrow{\dots} x_{p_1-1} \xleftarrow{\alpha_{p_1}} x \xrightarrow{\alpha'_{q_1}} x'_{q_1-1} \xrightarrow{\dots} x'_1 \xrightarrow{\alpha'_1} v, \\
 \Delta^{(2)} &= \bullet \xrightarrow{\beta_{p_2}} y_{p_2-1} \xrightarrow{\dots} y_1 \xrightarrow{\beta_1} y \xleftarrow{\beta'_1} y'_1 \xleftarrow{\dots} y'_{q_2-1} \xleftarrow{\beta'_{q_2}} v,
 \end{aligned}$$

and

$$\Delta^{(3)} = \bullet \xrightarrow{\gamma_{p_3}} z_{p_3-1} \xrightarrow{\dots} z_1 \xrightarrow{\gamma_1} z \xleftarrow{\gamma'_1} z'_1 \xleftarrow{\dots} z'_{q_3-1} \xleftarrow{\gamma'_{q_3}} v,$$

for some $p_1, p_2, p_3, q_1, q_2, q_3 \in \mathbb{N}_+$. Moreover, we may assume that $\beta_{p_2}\alpha_1 \in R$. Consequently, by shifting relations we may assume that $\beta_i\beta_{i+1} \in R$ for all $i \in [1, p_2 - 1]$. There are two cases we have to consider.

Assume first $\gamma_i\gamma_{i+1} \notin R$ for all $i \in [1, p_3 - 1]$. If (Δ', R') is the bound quiver obtained from (Δ, R) by applying the generalized APR-coreflection at u followed by the composition of the APR-coreflection at y_i and the generalized APR-coreflection at u for $i = p_2 - 1, \dots, 1$, then

$$R' = \begin{cases} R \setminus \{\beta_{p_2}\alpha_1, \beta_1\beta_2\} \cup \{\gamma_{p_3}\alpha_1, \beta_{p_2}\gamma_{p_3}\} & p_2 > 1 \\ R \setminus \{\beta_{p_2}\alpha_1\} \cup \{\gamma_{p_3}\alpha_1\} & p_2 = 1, \end{cases}$$

$$\Delta'^{(1)} = \bullet \xleftarrow{\alpha_1} z_{p_3-1} \xleftarrow{\dots} x_1 \xleftarrow{\alpha_{p_1}} x_{p_1-1} \xleftarrow{\alpha_{p_1}} x \xrightarrow{\alpha'_{q_1}} x'_{q_1-1} \xrightarrow{\dots} x'_1 \xrightarrow{\alpha'_1} v,$$

$$\Delta'(2) = \begin{array}{c} \bullet \\ z_{p_3-1} \end{array} \xrightarrow{\gamma_{p_3}} \begin{array}{c} \bullet \\ y_{p_2-1} \end{array} \xrightarrow{\beta_{p_2}} \begin{array}{c} \bullet \\ y_{p_2-2} \end{array} \cdots \rightarrow \begin{array}{c} \bullet \\ u \end{array} \xleftarrow{\beta_1} \begin{array}{c} \bullet \\ y \end{array} \xleftarrow{\beta'_1} \begin{array}{c} \bullet \\ y'_1 \end{array} \cdots \leftarrow \begin{array}{c} \bullet \\ y'_{q_2-1} \end{array} \xleftarrow{\beta'_{q_2}} \begin{array}{c} \bullet \\ v \end{array},$$

and

$$\Delta'(3) = \begin{array}{c} \bullet \\ z_{p_3-1} \end{array} \cdots \rightarrow \begin{array}{c} \bullet \\ z_1 \end{array} \xrightarrow{\gamma_1} \begin{array}{c} \bullet \\ z \end{array} \xleftarrow{\gamma'_1} \begin{array}{c} \bullet \\ z'_1 \end{array} \cdots \leftarrow \begin{array}{c} \bullet \\ z'_{q_3-1} \end{array} \xleftarrow{\gamma'_{q_3}} \begin{array}{c} \bullet \\ v \end{array},$$

where $z_{p_3-1} = z$ if $p_3 = 1$. Consequently, the claim follows by an easy induction.

Assume now that there exists $i \in [1, p_3 - 1]$ such that $\gamma_i \gamma_{i+1} \in R$. Consequently $p_2 = 1$. Moreover, by shifting relations we may assume that $\gamma_{p_3-1} \gamma_{p_3} \in R$. If (Δ', R') is the bound quiver obtained from (Δ, R) by applying the generalized APR-coreflection at u , then $R' = R \setminus \{\beta_1 \alpha_1, \gamma_{p_3-1} \gamma_{p_3}\} \cup \{\gamma_{p_3} \alpha_1, \gamma_{p_3-1} \beta_1\}$,

$$\Delta'(1) = \begin{array}{c} \bullet \\ u \end{array} \xleftarrow{\gamma_{p_3}} \begin{array}{c} \bullet \\ z_{p_3-1} \end{array} \xleftarrow{\alpha_1} \begin{array}{c} \bullet \\ x_1 \end{array} \cdots \leftarrow \begin{array}{c} \bullet \\ x_{p_1-1} \end{array} \xleftarrow{\alpha_{p_1}} \begin{array}{c} \bullet \\ x \end{array} \xrightarrow{\alpha'_{q_1}} \begin{array}{c} \bullet \\ x'_{q_1-1} \end{array} \cdots \rightarrow \begin{array}{c} \bullet \\ x'_1 \end{array} \xrightarrow{\alpha'_1} \begin{array}{c} \bullet \\ v \end{array},$$

$$\Delta'(2) = \begin{array}{c} \bullet \\ u \end{array} \xleftarrow{\beta_1} \begin{array}{c} \bullet \\ y \end{array} \xleftarrow{\beta'_1} \begin{array}{c} \bullet \\ y'_1 \end{array} \cdots \leftarrow \begin{array}{c} \bullet \\ y'_{q_2-1} \end{array} \xleftarrow{\beta'_{q_2}} \begin{array}{c} \bullet \\ u \end{array},$$

and

$$\Delta'(3) = \begin{array}{c} \bullet \\ u \end{array} \xrightarrow{\gamma_{p_3-1}} \begin{array}{c} \bullet \\ z_{p_3-2} \end{array} \cdots \rightarrow \begin{array}{c} \bullet \\ z \end{array} \xleftarrow{\gamma'_1} \begin{array}{c} \bullet \\ z'_1 \end{array} \cdots \leftarrow \begin{array}{c} \bullet \\ z'_{q_3-1} \end{array} \xleftarrow{\gamma'_{q_3}} \begin{array}{c} \bullet \\ u \end{array},$$

thus the claim follows. \square

Step 12. *We may assume that there is at most one $j \in [1, 3]$ such that $\Delta^{(j)}$ is not equioriented.*

Proof. If the above condition is not satisfied, then without loss of generality we may assume that

$$\Delta(1) = \begin{array}{c} \bullet \\ u \end{array} \xleftarrow{\alpha_1} \begin{array}{c} \bullet \\ x_1 \end{array} \cdots \leftarrow \begin{array}{c} \bullet \\ x_{p_1-1} \end{array} \xleftarrow{\alpha_{p_1}} \begin{array}{c} \bullet \\ x \end{array} \xrightarrow{\alpha'_{q_1}} \begin{array}{c} \bullet \\ x'_{q_1-1} \end{array} \cdots \rightarrow \begin{array}{c} \bullet \\ x'_1 \end{array} \xrightarrow{\alpha'_1} \begin{array}{c} \bullet \\ v \end{array},$$

$$\Delta(2) = \begin{array}{c} \bullet \\ u \end{array} \xrightarrow{\beta_{p_2}} \begin{array}{c} \bullet \\ y_{p_2-1} \end{array} \cdots \rightarrow \begin{array}{c} \bullet \\ y_1 \end{array} \xrightarrow{\beta_1} \begin{array}{c} \bullet \\ y \end{array} \xleftarrow{\beta'_1} \begin{array}{c} \bullet \\ y'_1 \end{array} \cdots \leftarrow \begin{array}{c} \bullet \\ y'_{q_2-1} \end{array} \xleftarrow{\beta'_{q_2}} \begin{array}{c} \bullet \\ v \end{array},$$

and

$$\Delta(3) = \begin{array}{c} \bullet \\ u \end{array} \xrightarrow{\gamma_{p_3}} \begin{array}{c} \bullet \\ z_{p_3-1} \end{array} \cdots \rightarrow \begin{array}{c} \bullet \\ z_1 \end{array} \xrightarrow{\gamma_1} \begin{array}{c} \bullet \\ v \end{array},$$

for some $p_1, p_2, p_3, q_1, q_2 \in \mathbb{N}_+$. In this proof we will again number the cases. Up to symmetry, there are three main cases we have to consider: either $\beta_{p_2} \alpha_1 \in R$ and $\beta'_{q_2} \alpha'_1 \in R$, or $\beta_{p_2} \alpha_1 \in R$ and $\beta'_{q_2} \gamma_1 \in R$, or $\gamma_{p_3} \alpha_1 \in R$ and $\beta'_{q_2} \gamma_1 \in R$.

(1) Assume $\beta_{p_2}\alpha_1 \in R$ and $\beta'_{q_2}\alpha'_1 \in R$. In this case we may apply the same arguments as in the proof of the previous step. Note however that if $\gamma_i\gamma_{i+1} \notin R$ for all $i \in [1, p_3 - 1]$, then we obtain a gentle bound quiver whose bound quiver algebra is tilting-cotilting equivalent to $\Lambda'_0(p, r)$ for some $p \in \mathbb{N}_+$ and $r \in [0, p - 1]$ according to Proposition 1.8.

(2) Assume that $\beta_{p_2}\alpha_1 \in R$ and $\beta'_{q_2}\gamma_1 \in R$.

(2.1) Assume that $\alpha'_i\alpha'_{i+1} \notin R$ for all $i \in [1, q_1 - 1]$ and $\beta'_i\beta'_{i+1} \notin R$ for all $i \in [1, q_2 - 1]$. By shifting the relation $\beta'_{q_2}\gamma_1$ to the left we may assume that $q_2 = 1$.

(2.1.1) Assume that $\beta_i\beta_{i+1} \notin R$ for all $i \in [1, p_2 - 1]$. By shifting the relation $\beta_{p_2}\alpha_1$ to the left we may assume that $p_2 = 1$. Consequently, the path algebra of the bound quiver obtained from (Δ, R) by application of the APR-reflections at $y, v, x'_1, \dots, x'_{q_1-1}$ is easily seen to be tilting-cotilting equivalent to $\Lambda_2(p_3+1, p_1+1, q_1, r_3, r_1)$, where r_1 is the number of $i \in [1, p_1 - 1]$ such that $\alpha_i\alpha_{i+1} \in R$ and r_3 is the number of $i \in [1, p_3 - 1]$ such that $\gamma_i\gamma_{i+1} \in R$.

(2.1.2) Assume that there exists $i \in [1, p_2 - 1]$ such that $\beta_i\beta_{i+1} \in R$. By shifting relations we may assume that $\beta_1\beta_2 \in R$. If (Δ', R') is the bound quiver obtained from (Δ, R) by applying the APR-reflections at $y, y_1, v, x'_1, \dots, x'_{q_1-1}$, then $R' = R \setminus \{\beta_1\beta_2, \beta'_1\gamma_1\} \cup \{\alpha'_{q_1}\beta_2\}$,

$$\Delta'^{(1)} = \bullet \xleftarrow{\alpha_1} \bullet \xleftarrow{\dots} \bullet \xleftarrow{\alpha_{p_1}} \bullet \xleftarrow{\alpha'_{q_1}} \bullet, \quad \begin{matrix} u \\ x_1 \\ x_{p_1-1} \\ x \\ x'_{q_1-1} \end{matrix}$$

$$\Delta'^{(2)} = \bullet \xrightarrow{\dots} \bullet \xrightarrow{\beta_2} \bullet, \quad \begin{matrix} u \\ y_2 \\ x'_{q_1-1} \end{matrix}$$

and

$$\Delta'^{(3)} = \bullet \xrightarrow{\gamma_{p_3}} \bullet \xrightarrow{\dots} \bullet \xrightarrow{\gamma_1} \bullet \xrightarrow{\beta_1} \bullet \xleftarrow{\beta'_1} \bullet \xleftarrow{\alpha'_1} \bullet \xleftarrow{\dots} \bullet, \quad \begin{matrix} u \\ z_{p_3-1} \\ z_1 \\ y_1 \\ y \\ v \\ x'_1 \\ x'_{q_1-1} \end{matrix}$$

where $x'_{q_1-1} = v$ if $q_1 = 1$, hence the claim follows.

(2.2) Assume that there exists $i \in [1, q_2 - 1]$ such that $\beta'_i\beta'_{i+1} \in R$. By shifting relations we may assume that $\beta'_1\beta'_2 \in R$. Moreover, this condition implies that $\beta_i\beta_{i+1} \notin R$ for all $i \in [1, p_2 - 1]$. By shifting the relation $\beta_{p_2}\alpha_1$ to the left we may assume $p_2 = 1$. If (Δ', R') is the bound quiver obtained from (Δ, R) by applying the APR-reflections at y and y'_1 , then $R' = R \setminus \{\beta_1\alpha_1, \beta'_1\beta'_2\} \cup \{\beta_1\beta'_2\}$,

$$\Delta'^{(1)} = \bullet \xleftarrow{\beta'_1} \bullet \xleftarrow{\alpha_1} \bullet \xleftarrow{\dots} \bullet \xleftarrow{\alpha_{p_1}} \bullet \xrightarrow{\alpha'_{q_1}} \bullet \xrightarrow{\dots} \bullet \xrightarrow{\alpha'_1} \bullet, \quad \begin{matrix} y \\ y'_1 \\ x_1 \\ x_{p_1-1} \\ x \\ x'_{q_1-1} \\ x'_1 \\ v \end{matrix}$$

$$\Delta'^{(2)} = \bullet \xleftarrow{\beta'_2} \bullet \xleftarrow{\dots} \bullet, \quad \begin{matrix} y \\ y'_2 \\ v \end{matrix}$$

and

$$\Delta'^{(3)} = \bullet_y \xrightarrow{\beta_1} \bullet_u \xrightarrow{\gamma_{p_3}} \bullet_{z_{p_3-1}} \cdots \xrightarrow{\gamma_1} \bullet_{z_1} \xrightarrow{\gamma_1} \bullet_v,$$

hence the claim follows again.

(2.3) Assume that $\beta'_i \beta'_{i+1} \notin R$ for all $i \in [1, q_2 - 1]$ and there exists $i \in [1, q_1 - 1]$ such that $\alpha'_i \alpha'_{i+1} \in R$. Observe that in this case $\alpha_i \alpha_{i+1} \notin R$ for all $i \in [1, p_1 - 1]$, hence by shifting the relation $\beta_{p_2} \alpha_1$ to the right we may assume that $p_1 = 1$. Similarly, $\gamma_i \gamma_{i+1} \notin R$ for all $i \in [1, p_3 - 1]$.

(2.3.1) Assume that there exists $i \in [1, p_2 - 1]$ such that $\beta_i \beta_{i+1} \in R$, then by shifting relations we may assume that $\beta_1 \beta_2 \in R$. Moreover, by shifting the relation $\beta'_{q_2} \gamma_1$ to the left we may assume that $q_2 = 1$. Additionally, by shifting relations we may assume that $\alpha'_1 \alpha'_2 \in R$. If (Δ', R') is the bound quiver obtained from (Δ, R) by applying the HW-reflection at y followed by the APR-coreflection at y and the APR-reflections at v, z_1, \dots, z_{p_3-1} , then $R' = R \setminus \{\alpha'_1 \alpha'_2, \beta_1 \beta_2, \beta'_1 \gamma_1\} \cup \{\gamma_{p_3} \alpha'_2\}$,

$$\Delta'^{(1)} = \bullet_u \xleftarrow{\alpha_1} \bullet_x \cdots \xrightarrow{\alpha'_2} \bullet_{x'_2} \xrightarrow{\alpha'_2} \bullet_{z_{p_3-1}},$$

$$\Delta'^{(2)} = \bullet_u \xrightarrow{\beta_{p_2}} \bullet_{y_{p_2-1}} \cdots \xrightarrow{\beta_1} \bullet_{y_1} \xrightarrow{\beta_1} \bullet_y \xleftarrow{\beta'_1} \bullet_{x'_1} \xleftarrow{\alpha'_1} \bullet_v \xleftarrow{\gamma_1} \bullet_{z_1} \cdots \xrightarrow{\gamma_{p_3-1}} \bullet_{z_{p_3-1}},$$

and

$$\Delta'^{(3)} = \bullet_u \xleftarrow{\gamma_{p_3}} \bullet_{z_{p_3-1}},$$

where $z_{p_3-1} = v$ if $p_3 = 1$. In particular, we reduce the proof to the situation dual either to (2.1) or to (2.2).

(2.3.2) Assume that $\beta_i \beta_{i+1} \notin R$ for all $i \in [1, p_2 - 1]$. By shifting the relation $\beta'_{q_2} \gamma_1$ to the right we may assume that $p_3 = 1$. Additionally, by shifting relations we may assume that $\alpha'_{q_1-1} \alpha'_{q_1} \in R$. The bound quiver algebra of the bound quiver obtained from (Δ, R) by applying the APR-coreflections at $x, x'_{q_1-1}, u, x, x'_{q_1-1}, y_{p_2-1}, \dots, y_1$ is easily seen to be tilting-cotilting equivalent to $\Lambda_2(q_2 + 1, q_1, p_2 + 1, 0, r'_1 - 1)$, where r'_1 is the number of $i \in [1, q_1 - 1]$ such that $\alpha'_i \alpha'_{i+1} \in R$.

(3) Assume that $\gamma_{p_3} \alpha_1 \in R$ and $\beta'_{q_2} \gamma_1 \in R$.

(3.1) Assume that there exists $i \in [1, p_2 - 1]$ such that $\beta_i \beta_{i+1} \in R$. By shifting relations we may assume that $\beta_{p_2-1} \beta_{p_2} \in R$. Since in this case $\gamma_i \gamma_{i+1} \notin R$ for all $i \in [1, p_3 - 1]$, we may assume, by shifting the relation $\gamma_{p_3} \alpha_1$ to the left, that $p_3 = 1$. Consequently, the bound quiver algebra of the bound quiver obtained from (Δ, R) by applying the generalized APR-coreflection at u is tilting-cotilting equivalent to $\Lambda'_0(p, r)$ for some $p \in \mathbb{N}_+$ and $r \in [0, p - 1]$ according to Proposition 1.8.

(3.2) Assume that $\beta_i \beta_{i+1} \notin R$ for all $i \in [1, p_2 - 1]$. By shifting relations we may also assume that $\gamma_i \gamma_{i+1} \in R$ for all $i \in [1, p_3 - 1]$. If

(Δ', R') is the bound quiver obtained from (Δ, R) by applying the APR-coreflection at u followed by the composition of the APR-coreflection at z_i and the generalized APR-coreflection at u for $i = p_3 - 1, \dots, 1$, then

$$R' = \begin{cases} R \setminus \{\gamma_{p_3}\alpha_1, \gamma_1\gamma_2, \beta'_{q_2}\gamma_1\} \cup \{\beta_{p_2}\alpha_1, \gamma_{p_3}\beta_{p_2}, \beta'_{q_2}\gamma_2\} & p_3 > 1, \\ R \setminus \{\gamma_1\alpha_1, \beta'_{q_2}\gamma_1\} \cup \{\beta_{p_2}\alpha_1, \beta'_{q_2}\beta_{p_2}\} & p_3 = 1, \end{cases}$$

$$\Delta'^{(1)} = \begin{array}{ccccccccccc} \bullet & \xleftarrow{\alpha_1} & \bullet & \xleftarrow{\dots} & \bullet & \xleftarrow{\alpha_{p_1}} & \bullet & \xrightarrow{\alpha'_{q_1}} & \bullet & \xrightarrow{\dots} & \bullet & \xrightarrow{\alpha'_1} & \bullet & \xrightarrow{\gamma_{p_3}} & \bullet \\ y_{p_2-1} & & x_1 & & & & x_{p_1-1} & & x & & x'_{q_1-1} & & x'_1 & & v & & u \end{array},$$


$$\Delta'^{(2)} = \begin{array}{ccccccccccc} \bullet & \xrightarrow{\dots} & \bullet & \xrightarrow{\beta_1} & \bullet & \xleftarrow{\beta'_1} & \bullet & \xleftarrow{\dots} & \bullet & \xleftarrow{\beta'_{q_2}} & \bullet \\ y_{p_2-1} & & & & y_1 & & y & & y'_1 & & y'_{q_2-1} & & & & u \end{array}$$

and

$$\Delta'^{(3)} = \begin{array}{ccccccc} \bullet & \xrightarrow{\beta_{p_2}} & \bullet & \xrightarrow{\dots} & \bullet & \xrightarrow{\gamma_2} & \bullet \\ y_{p_2-1} & & z_{p_3-1} & & & & z_1 & & u \end{array},$$

where $y_{p_2-1} = y$ if $p_2 = 1$. Consequently, the claim follows by induction. \square

For $p_1, p_2, p_3 \in \mathbb{N}_+$, $p_2 \geq 2$, $r_1 \in [0, p_1 - 1]$, and $r_2 \in [1, p_2 - 1]$, let $\Lambda'_2(p_1, p_2, p_3, r_1, r_2)$ be the algebra of the quiver



bound by $\alpha_i\alpha_{i+1}$ for $i \in [p_1 - r_1, p_1 - 1]$, $\alpha_{p_1}\beta$, $\beta\alpha_1$, $\gamma_i\gamma_{i+1}$ for $i \in [1, r_2]$. Observe that $\Lambda'_2(p_1, p_2, p_3, r_1, r_2)$ is tilting-cotilting equivalent to $\Lambda_2(p_2, p_1 + 1, p_3, r_2 - 1, r_1 + 1)$. Indeed, it is enough to apply the HW-reflection at x_i followed by the APR-coreflection at x_i for $i = 1, \dots, p_3$, where $x_i = t\delta_i$ for $i \in [1, p_3]$.

Step 13. We may assume that $\Delta^{(j)}$ is equioriented for each $j \in [1, 3]$.

Proof. Suppose that there exists $j \in [1, 3]$ such that $\Delta^{(j)}$ is not equioriented. Without loss of generality we may assume that

$$\Delta^{(1)} = \begin{array}{ccccccccccc} \bullet & \xleftarrow{\alpha_1} & \bullet & \xleftarrow{\dots} & \bullet & \xleftarrow{\alpha_{p_1}} & \bullet & \xrightarrow{\alpha'_{q_1}} & \bullet & \xrightarrow{\dots} & \bullet & \xrightarrow{\alpha'_1} & \bullet \\ u & & x_1 & & & & x_{p_1-1} & & x & & x'_{q_1-1} & & x'_1 & & v \end{array},$$

$$\Delta^{(2)} = \begin{array}{ccccccc} \bullet & \xrightarrow{\beta_{p_2}} & \bullet & \xrightarrow{\dots} & \bullet & \xrightarrow{\beta_1} & \bullet \\ u & & y_{p_2-1} & & & & y_1 & & v \end{array},$$

and

$$\Delta^{(3)} = \bullet \xleftarrow{\gamma_1} z_1 \xleftarrow{\dots} z_{p_3-1} \xleftarrow{\gamma_{p_3}} v ,$$

for some $p_1, p_2, p_3, q_1 \in \mathbb{N}_+$. We may additionally assume that $\alpha_{i+1}\alpha_i \notin R$ for all $i \in [1, p_1 - 1]$. Let r'_1 be the number of $i \in [1, q_1 - 1]$ such that $\alpha'_i\alpha'_{i+1} \in R$, let r_2 be the number of $i \in [1, p_2 - 1]$ such that $\beta_i\beta_{i+1} \in R$, and let r_3 be the number of $i \in [1, p_3 - 1]$ such that $\gamma_i\gamma_{i+1} \in R$. Observe that by symmetry we may assume that $r'_1 > 0$ if $\gamma_{p_3}\alpha'_1 \in R$ and $\beta_{p_2}\alpha_1 \notin R$.

Assume first that $\beta_{p_2}\alpha_1 \in R$. In this case by shifting the relation $\beta_{p_2}\alpha_1$ to the right we may assume that $p_1 = 1$. Observe that either $r'_1 = 0$ or $r_2 = 0$.

If $r_3 \geq 1$, then by shifting relations we may assume that $\gamma_1\gamma_2 \in R$. If (Δ', R') is the bound quiver obtained from (Δ, R) by applying the generalized APR-reflection at u , then $R' = R \setminus \{\beta_{p_2}\alpha_1, \gamma_1\gamma_2\} \cup \{\alpha_1\gamma_2, \beta_{p_2}\gamma_1\}$,

$$\Delta'^{(1)} = \bullet \xrightarrow{\alpha_1} x \xrightarrow{\alpha'_{q_1}} x'_{q_1-1} \xrightarrow{\dots} x'_1 \xrightarrow{\alpha'_1} v ,$$

$$\Delta'^{(2)} = \bullet \xrightarrow{\gamma_1} z_1 \xrightarrow{\beta_{p_2}} y_{p_2-1} \xrightarrow{\dots} y_1 \xrightarrow{\beta_1} v ,$$

and

$$\Delta'^{(3)} = \bullet \xleftarrow{\gamma_2} z_2 \xleftarrow{\dots} v ,$$

hence the claim follows in this case.

Assume now that $r_3 = 0$. There are two additional possibilities in this case. If $\gamma_{p_3}\alpha'_1 \in R$, then $r_2 \geq 1$ (since (Δ, R) is a bound quiver). Consequently, $r'_1 = 0$ and we have the situation symmetric to the previous one. On the other hand, if $\gamma_{p_3}\beta_1 \in R$, then by shifting the relation $\gamma_{p_3}\beta_1$ to the left we may assume that $p_3 = 1$. Consequently, if (Δ', R') is the bound quiver obtained from (Δ, R) by applying the generalized APR-reflection at u , then $R' = R \setminus \{\beta_{p_2}\alpha_1, \gamma_1\beta_1\} \cup \{\alpha_1\beta_1, \beta_{p_2}\gamma_1\}$,

$$\Delta'^{(1)} = \bullet \xrightarrow{\alpha_1} x \xrightarrow{\alpha'_{q_1}} x'_{q_1-1} \xrightarrow{\dots} x'_1 \xrightarrow{\alpha'_1} v ,$$

$$\Delta'^{(2)} = \bullet \xleftarrow{\beta_1} y_1 \xleftarrow{\dots} y_{p_2-1} \xleftarrow{\beta_{p_2}} v ,$$

and

$$\Delta'^{(3)} = \bullet \xrightarrow{\gamma_1} v ,$$

hence the claim follows.

Assume now that $\beta_{p_2}\gamma_1 \in R$. If $\gamma_{p_3}\beta_1 \in R$, then it follows easily that the bound quiver algebra of (Δ, R) is tilting-cotilting equivalent either to $\Lambda_2(p_2 + p_3 - r_2 - 1, r_2 + 1, q_1, p_1, r_3)^{\text{op}}$ if $r'_1 = 0$, or to $\Lambda'_2(p_2 + p_3 - 1, q_1, p_1, r_3, r'_1)^{\text{op}}$ if $r'_1 \geq 1$. Since $\Lambda'_2(p_2 + p_3 - 1, q_1, p_1, r_3, r'_1)$ is tilting-cotilting equivalent to $\Lambda_2(q_1, p_2 + p_3, p_1, r'_1 - 1, r_3 + 1)$, thus we may assume that $\gamma_{p_3}\alpha'_1 \in R$. Consequently, by shifting relations we may assume that $\alpha'_i\alpha'_{i+1} \in R$ for all $i \in [1, q_1 - 1]$. Recall that $q_1 > 1$ in this case. If (Δ', R') is the bound quiver obtained from (Δ, R) by applying the APR-coreflection at x followed by the composition of the HW-coreflection at x'_{q_1-1} and the APR-reflection at x'_{q_1-1} applied $q_1 - 1$ times, then $R' = R \setminus \{\alpha'_{q_1-1}\alpha'_{q_1}\} \cup \{\alpha'_{q_1-1}\alpha_{p_1}\}$,

$$\Delta'(1) = \begin{cases} \bullet \xleftarrow{\alpha_1} \bullet \xleftarrow{\dots} \bullet \xrightarrow{\alpha_{p_1}} \bullet \xrightarrow{\alpha'_{q_1-1}} \bullet \xrightarrow{\dots} \bullet & p_1 > 1, \\ \bullet \xrightarrow{\alpha_1} \bullet \xrightarrow{\alpha'_{q_1-1}} \bullet \xrightarrow{\dots} \bullet & p_1 = 1, \end{cases}$$

$$\Delta'(2) = \bullet \xrightarrow{\beta_1} \bullet \xrightarrow{\dots} \bullet \xrightarrow{\beta_{p_2}} \bullet \xrightarrow{\alpha'_{q_1}} \bullet,$$

and

$$\Delta'(3) = \bullet \xleftarrow{\gamma_1} \bullet \xleftarrow{\dots} \bullet \xleftarrow{\gamma_{p_3}} \bullet,$$

thus the claim follows by induction. \square

We may prove now Proposition 2.1. According to our considerations we may assume that (Δ, R) is proper,

$$\Delta(1) = \bullet \xleftarrow{\alpha_1} \bullet \xleftarrow{\dots} \bullet \xleftarrow{\alpha_{p_1}} \bullet,$$

$$\Delta(2) = \bullet \xrightarrow{\beta_{p_2}} \bullet \xrightarrow{\dots} \bullet \xrightarrow{\beta_1} \bullet,$$

and

$$\Delta(3) = \bullet \xleftarrow{\gamma_1} \bullet \xleftarrow{\dots} \bullet \xleftarrow{\gamma_{p_3}} \bullet,$$

for some $p_1, p_2, p_3 \in \mathbb{N}_+$. Moreover, we may additionally assume that $\alpha_i\alpha_{i+1} \notin R$ for all $i \in [1, p_1 - 1]$. Let r_2 be the number of $i \in [1, p_2 - 1]$ such that $\beta_i\beta_{i+1} \in R$ and let r_3 be the number of $i \in [1, p_3 - 1]$ such that $\gamma_i\gamma_{i+1} \in R$.

Observe that if either $\beta_{p_2}\alpha_1 \in R$ and $\gamma_{p_3}\beta_1 \in R$ or $\beta_{p_2}\gamma_1 \in R$ and $\alpha_{p_1}\beta_1 \in R$, then the claim follows from Proposition 1.16, thus we have to consider two remaining cases.

Assume first that $\beta_{p_2}\alpha_1 \in R$ and $\alpha_{p_1}\beta_1 \in R$. In this case by shifting the relation $\beta_{p_2}\alpha_1$ to the right we may assume that $p_1 = 1$. If $r_3 = 0$,

then the bound quiver algebra of (Δ, R) is tilting-cotilting equivalent to the $\Lambda_1(p_2, 1, p_3, 0, r_2)$ (observe that $r_2 \geq 1$ since (Δ, R) is a bound quiver). On the other hand, if there exists $i \in [1, p_3 - 1]$ such that $\gamma_i \gamma_{i+1} \in R$, then by shifting relations we may assume that $\gamma_1 \gamma_2 \in R$. Consequently, the bound quiver algebra of the bound quiver obtained from (Δ, R) by applying the generalized APR-reflection at u is tilting-cotilting equivalent to $\Lambda_2(p_2 + 1, p_3, 0, r_2 + 1, r_3 - 1)$ and this finishes the proof in this case.

Assume now that $\beta_{p_2} \gamma_1 \in R$ and $\gamma_{p_3} \beta_1 \in R$. In this case it follows by shifting relations that the bound quiver algebra of (Δ, R) is tilting-cotilting equivalent to $\Lambda_1(p_2 + p_3 - r_3 - 1, r_3 + 1, p_1, 0, r_2)$ (again $r_2 \geq 1$ since (Δ, R) is a bound quiver) and this finishes the proof.

3. MINIMALITY OF THE LIST

In this section we prove that different algebras from the list in Theorem 1 are not derived equivalent. We also check that the algebras listed in Theorem 1 are nondegenerate, while the algebras listed in Theorem 2 are degenerate. A tool used in order to distinguish between derived equivalence classes of these algebras will be the derived invariant introduced by Avella-Alaminos and Geiss in [7].

Let (Δ, R) be a gentle quiver. By a permitted thread in (Δ, R) we mean either a maximal path in (Δ, R) or $x \in \Delta_0$ such that there is at most one arrow α with $s\alpha = x$, there is at most one arrow β with $t\beta = x$, and $\alpha\beta \notin R$ for all $\alpha, \beta \in \Delta_1$ with $s\alpha = x = t\beta$. Similarly, we define notion of a forbidden thread in (Δ, R) . Namely, first we say that by an anti-path in (Δ, R) we mean a path $\alpha_1 \cdots \alpha_n$ in Δ such that $\alpha_i \alpha_{i+1} \in R$ for all $i \in [1, n - 1]$. In particular, every trivial path is an anti-path. By a forbidden thread we mean either a maximal anti-path in (Δ, R) or $x \in \Delta_0$ such that there is at most one arrow α with $s\alpha = x$, there is at most one arrow β with $t\beta = x$, and $\alpha\beta \in R$ for all $\alpha, \beta \in \Delta_1$ with $s\alpha = x = t\beta$.

By a characteristic sequence in a gentle bound quiver (Δ, R) we mean a sequence $(\sigma_i, \tau_i)_{i \in \mathbb{Z}}$ of permitted threads σ_i , $i \in \mathbb{Z}$, and forbidden threads τ_i , $i \in \mathbb{Z}$, such that for each $i \in \mathbb{Z}$ the following conditions are satisfied:

- (1) $t\tau_i = t\sigma_i$ and $s\sigma_{i+1} = s\tau_i$,
- (2) if $\sigma_i = x = \tau_i$ for $x \in \Delta_0$ then $\sigma_{i+1} \neq x$, unless $\Delta_1 = \emptyset$,
- (3) if $\tau_i = x = \sigma_{i+1}$ for $x \in \Delta_0$, then $\tau_{i+1} \neq x$, unless $\Delta_1 = \emptyset$,
- (4) if neither σ_i nor τ_i is a trivial path, then the terminating arrow of τ_i differs from the terminating arrow of σ_i ,
- (5) if neither τ_i nor σ_{i+1} is a trivial path, then the starting arrow of σ_{i+1} differs from the starting arrow of τ_i .

We identify characteristic sequences $(\sigma_i, \tau_i)_{i \in \mathbb{Z}}$ and $(\sigma'_i, \tau'_i)_{i \in \mathbb{Z}}$ if there exists $l \in \mathbb{Z}$ such that $\sigma_i = \sigma'_{i+l}$ and $\tau_i = \tau'_{i+l}$ for all $i \in \mathbb{Z}$. By the type

of the characteristic sequence $(\sigma_i, \tau_i)_{i \in \mathbb{Z}}$ we mean a pair $(n, m) \in \mathbb{N} \times \mathbb{N}$ defined by $n = \min\{l \in \mathbb{N}_+ \mid \sigma_l = \sigma_0\}$ and $m = \sum_{i \in [1, n]} \ell(\tau_i)$. In the above situation we also write $(\sigma_1, \tau_1, \dots, \sigma_n, \tau_n)$ instead of $(\sigma_i, \tau_i)_{i \in \mathbb{Z}}$. Additionally, we also call a sequence $(\alpha_i)_{i \in \mathbb{Z}}$ of arrows in Δ a characteristic sequence if $s\alpha_i = t\alpha_{i+1}$ and $\alpha_i\alpha_{i+1} \in R$ for all $i \in \mathbb{Z}$. Again we identify sequences $(\alpha_i)_{i \in \mathbb{Z}}$ and $(\alpha'_i)_{i \in \mathbb{Z}}$ if there exists $l \in \mathbb{Z}$ such that $\alpha_i = \alpha'_{i+l}$ for all $i \in \mathbb{Z}$. The type of a characteristic sequence $(\alpha_i)_{i \in \mathbb{Z}}$ of the above type is by definition $(0, m)$, where $m = \min\{l \in \mathbb{N}_+ \mid \alpha_l = \alpha_0\}$. In the above situation we also write $\alpha_1 \cdots \alpha_m$ instead of $(\alpha_i)_{i \in \mathbb{Z}}$.

If (Δ, R) is a gentle bound quiver, then the function $\phi_{\Delta, R} : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$, where $\phi_{\Delta, R}(n, m)$ is the number of the characteristic sequences of type (n, m) for $(n, m) \in \mathbb{N} \times \mathbb{N}$, is a derived invariant, i.e. if (Δ, R) and (Δ', R') are derived equivalent gentle bound quivers, then $\phi_{\Delta, R} = \phi_{\Delta', R'}$. If Λ is the bound quiver algebra of a gentle bound quiver (Δ, R) , then we also write ϕ_Λ instead of $\phi_{\Delta, R}$. We will write $\phi_{\Delta, R}$ as a ‘‘multi-set’’ $[(n_1, m_1), \dots, (n_l, m_l)]$, where (n, m) appears $\phi_{\Delta, R}(n, m)$ times.

We calculate the values of the above invariant for algebras appearing in Theorems 1 and 2, and this will finish the proofs of these theorems. The proof of the following lemma we leave to the reader as an easy exercise.

Lemma 3.1. *We have the following.*

(1) *If $p \in \mathbb{N}_+$ and $r \in [0, p - 1]$, then*

$$\phi_{\Lambda_0(p, r)} = [(p, p + 2)].$$

(2) *If $p \in \mathbb{N}_+$, then*

$$\phi_{\Lambda'_0(p, 0)} = [(p + 1, p + 3)].$$

(3) *If $p_1, p_2 \in \mathbb{N}_+$, $p_3, p_4 \in \mathbb{N}$, and $r_1 \in [0, p_1 - 1]$, are such that $p_2 + p_3 \geq 2$ and $r_1 + p_4 \geq 1$, then*

$$\phi_{\Lambda_1(p_1, p_2, p_3, p_4, r_1)} = [(p_1 - r_1 - 1, p_1 + p_2), (p_2 + p_3 - 1, p_3), (r_1 + p_4, p_4)].$$

(4) *If $p_1, p_2 \in \mathbb{N}_+$, $p_3 \in \mathbb{N}$, $r_1 \in [0, p_1 - 1]$, $r_2 \in [0, p_2 - 1]$, are such that $p_3 + r_1 + r_2 \geq 1$, then*

$$\phi_{\Lambda_2(p_1, p_2, p_3, r_1, r_2)} = [(p_1 - r_1 - 1, p_1), (p_2 - r_2 - 1, p_2), (r_1 + r_2 + p_3, p_3)].$$

REFERENCES

- [1] H. Asashiba, *The derived equivalence classification of representation-finite self-injective algebras*, J. Algebra **214** (1999), 182–221.
- [2] I. Assem and D. Happel, *Generalized tilted algebras of type A_n* , Comm. Algebra **9** (1981), 2101–2125.
- [3] I. Assem, D. Simson, and A. Skowroński, *Elements of the Representation Theory of Associative Algebras 1: Techniques of Representation Theory*, London Mathematical Society Student Texts, vol. 65, Cambridge University Press, Cambridge, 2006.
- [4] I. Assem and A. Skowroński, *Iterated tilted algebras of type \tilde{A}_n* , Math. Z. **195** (1987), 269–290.

- [5] ———, *Algebras with cycle-finite derived categories*, Math. Ann. **280** (1988), 441–463.
- [6] M. Auslander, M. I. Platzeck, and I. Reiten, *Coxeter functors without diagrams*, Trans. Amer. Math. Soc. **250** (1979), 1–46.
- [7] D. Avella-Alaminos and Ch. Geiss, *Combinatorial derived invariants for gentle algebras* (arXiv:math.RT/0607348).
- [8] M. Barot and H. Lenzing, *One-point extensions and derived equivalence*, J. Algebra **264** (2003), 1–5.
- [9] J. Białkowski, T. Holm, and A. Skowroński, *Derived equivalences for tame weakly symmetric algebras having only periodic modules*, J. Algebra **269** (2003), 652–668.
- [10] G. Bobiński, Ch. Geiss, and A. Skowroński, *Classification of discrete derived categories*, Cent. Eur. J. Math. **2** (2004), 19–49 (electronic).
- [11] R. Bocian, T. Holm, and A. Skowroński, *Derived equivalence classification of one-parametric self-injective algebras*, J. Pure Appl. Algebra **207** (2006), 491–536.
- [12] K. Bongartz, *Tilted algebras*, pp. 26–38.
- [13] S. Brenner and M. C. R. Butler, *Generalizations of the Bernstein-Gelfand-Ponomarev reflection functors*, 103–169.
- [14] T. Brüstle, *Derived-tame tree algebras*, Compositio Math. **129** (2001), 301–323.
- [15] Ch. Geiss, *Derived tame algebras and Euler-forms*, Math. Z. **239** (2002), 829–862, With an appendix by the author and B. Keller.
- [16] D. Happel, *On the derived category of a finite-dimensional algebra*, Comment. Math. Helv. **62** (1987), 339–389.
- [17] ———, *Triangulated Categories in the Representation Theory of Finite-Dimensional Algebras*, London Mathematical Society Lecture Note Series, vol. 119, Cambridge University Press, Cambridge, 1988.
- [18] ———, *Auslander-Reiten triangles in derived categories of finite-dimensional algebras*, Proc. Amer. Math. Soc. **112** (1991), 641–648.
- [19] D. Happel and C. M. Ringel, *Tilted algebras*, Trans. Amer. Math. Soc. **274** (1982), 399–443.
- [20] T. Holm, *Derived equivalence classification of algebras of dihedral, semidiheral, and quaternion type*, J. Algebra **211** (1999), 159–205.
- [21] D. Hughes and J. Waschbüsch, *Trivial extensions of tilted algebras*, Proc. London Math. Soc. (3) **46** (1983), 347–364.
- [22] B. Keller, *Deriving DG categories*, Ann. Sci. École Norm. Sup. (4) **27** (1994), 63–102.
- [23] Z. Pogorzały and A. Skowroński, *Self-injective biserial standard algebras*, J. Algebra **138** (1991), 491–504.
- [24] J. Rickard, *Morita theory for derived categories*, J. London Math. Soc. (2) **39** (1989), 436–456.
- [25] ———, *Derived categories and stable equivalence*, J. Pure Appl. Algebra **61** (1989), 303–317.
- [26] J. Schröer and A. Zimmermann, *Stable endomorphism algebras of modules over special biserial algebras*, Math. Z. **244** (2003), 515–530.
- [27] A. Skowroński and J. Waschbüsch, *Representation-finite biserial algebras*, J. Reine Angew. Math. **345** (1983), 172–181.
- [28] H. Tachikawa and T. Wakamatsu, *Applications of reflection functors for self-injective algebras*, 1986, pp. 308–327.
- [29] J.-L. Verdier, *Catégories dérivées*, pp. 262–311.
- [30] D. Vossieck, *The algebras with discrete derived category*, J. Algebra **243** (2001), 168–176.

FACULTY OF MATHEMATICS AND COMPUTER SCIENCE, NICOLAUS COPERNICUS UNIVERSITY, UL. CHOPINA 12/18, 87-100 TORUŃ, POLAND

E-mail address: `gregbob@mat.uni.torun.pl`

E-mail address: `pmalicki@mat.uni.torun.pl`