

EXTREME ANOMALIES OF WINTER AIR TEMPERATURE IN MID-LATITUDE EUROPE

J. OTTERMAN¹, J. ARDIZZONE², R. ATLAS³, D. BUNGATO²,
J. CIERNIEWSKI⁴, J. C. JUSEM², R. PRZYBYLAK⁵, S. SCHUBERT³, D. STARR³,
J. WALCZEWSKI⁶, A. WOS⁴

¹ Land-Atmosphere-Ocean-Research; at Data Assimilation Office, Code 910.3, NASA GSFC, Greenbelt, MD 20771, USA; ² General Science Corp., Beltsville, MD, USA; ³ Laboratory for Atmospheres, NASA GSFC, Greenbelt, MD 20771, USA; ⁴ Adam Mickiewicz University, Poznań, Poland; ⁵ Nicholas Copernicus University, Toruń, Poland; ⁶ Institute of Meteorology and Water Management, Branch of Cracow, Poland

As long as the earth remains, seed-time and harvest, and cold and heat, and summer and winter, and day and night, shall not cease

Genesis, Chapter VIII, Verse 22

ABSTRACT. The aim of this paper is to report extreme winter/early-spring air temperature (hereinafter temperature) anomalies in mid-latitude Europe, and to discuss the underlying forcing to these interannual fluctuations. Warm advection from the North Atlantic in late winter controls the surface-air temperature, as indicated by the substantial correlation between the speed of the surface southwesterlies over the eastern North Atlantic (quantified by a specific Index I_{na}) and the 2-meter level air temperatures (hereinafter T_s) over Europe, 45–60°N, in winter. In mid-March and subsequently, the correlation drops drastically (quite often it is negative). This change in the relationship between T_s and I_{na} marks a transition in the control of the surface-air temperature: absorption of insolation replaces the warm advection as the dominant control. This forcing by maritime-air advection in winter was demonstrated in a previous publication, and is re-examined here in conjunction with extreme fluctuations of temperatures in Europe. We analyze here the interannual variability at its extreme by comparing the warm-winter/early-spring of 1989/90 with the opposite scenario in 1995/96. For these two December-to-March periods the differences in the monthly mean air temperature in Warsaw and Toruń, Poland, range above 10°C. Short-term (shorter than a month) fluctuations of air temperature are likewise very strong. We conduct pentad-by-pentad analysis of the surface-maximum air temperature (hereinafter T_{max}), in a selected location, examining the dependence on I_{na} . The increased cloudiness and larger amounts of total precipitable water, corollary effects to the warm low-level advection in the 1989/90 winter, enhance the positive air temperature anomalies. The analysis of the ocean-surface winds is based on the Special Sensor Microwave/Imager (SSM/I) dataset; ascent rates, and over land wind data are from the European Centre for Medium-Range Weather Forecasts (ECMWF); maps of 2-m air temperature, cloud cover and precipitable water are from the National Centers for Environmental Prediction (NCEP) Reanalysis.

KEY WORDS: anomalies of air temperature in Europe, maritime-air advection, climatic fluctuations.

INTRODUCTION

The aim of this paper is to analyze the interannual variability which characterizes the late-winter surface air temperature in mid-latitude Europe, 45–60°N, and discuss its causes. Strong variability applies as well to the timing of the end of winter and onset of spring, whether measured by the beginning/end of snow cover, or by phenological events (Wos 1999; Jaagus and Ahas 2000). The fluctuations have profound implications for agriculture, forestry, and indeed for the way of life of the large population living in these regions.

With the winter sun low above the horizon, and low absorptivity (high albedo) of the snow-covered surface, advection from the oceans effectively controls the air temperatures over the adjoining continent, as shown by Rogers and Mosley-Thompson (1995), for instance. The surface of the North Atlantic is much warmer than that of the mid-latitude Europe, whereas the Arctic is colder, such that the air temperature in Europe fluctuate with the wind direction. It is widely recognized that circulation patterns over the North Atlantic and Europe fluctuate with the North Atlantic Oscillation (NAO), the stage of which is quantified by an Index (Rogers 1997). Recently, a specific index quantifying the low-level flow from the North Atlantic into Europe has been developed by measuring the strength of the ocean-surface southwesterlies over the eastern North Atlantic. Apparently, this index, I_{na} , is a more directly relevant measure of the maritime-air advection into France and into mid-latitude strips through Europe, from France to Russia (Otterman *et al.* 1999).

We analyze here the interannual variability at its extreme by comparing the warm-winter/early-spring of 1989/90 with the opposite scenario in 1995/96. Short-term (shorter than a month) fluctuations of the air temperature are likewise very strong, so for these two December-to-March periods we conduct pentad-by-pentad analysis of the T_{max} in a selected location and our Index I_{na} .

SSM/I, ECMWF, NCEP DATASETS

A large dataset of surface wind speeds over the global oceans has been derived from the SSM/I aboard the Defense Meteorological Satellite Program (DMSP) satellites. Variational analysis method was selected to derive the wind-vector data. In this method, the SSM/I retrievals (that is, the measurements of speed) are combined with independent wind observations to produce consistent fields of wind speed and direction. The resulting global ocean-surface dataset is appropriate for climate analysis (Atlas *et al.* 1996).

The ECMWF dataset is part of the Basic Level III-A analysis product with the ECMWF/World Climate Research Programme (WCRP) Global Atmospheric Data Archive. In our study, we use data interpolated horizontally to the standard

Goddard Laboratory for Atmospheres grid of 2.0° in latitude and 2.5° in longitude, from the ECMWF data archived on a 2.5° by 2.5° latitude/longitude grid.

Significant parts of our study are based on the NCEP Reanalysis dataset, described in detail by Kalnay et al. (1996), which extends from January 1948 essentially to the present. Improvements to the numerical weather prediction operational systems were introduced when satellite measurements become available (see Kalnay et al. 1996, for a documentation of the changes). The intent in reprocessing was to produce a consistent dataset. Still, some discontinuity was apparently introduced starting with 1979, relative to the earlier (1948–1978) period when no satellite observations were available. This uncertainty, crucially important to the evaluation of trends, is addressed in a recent report on the Reanalysis project (Kistler et al. 2001).

NORTH-ATLANTIC WINDS AS THE PRIMARY CONTROL OF THE SURFACE-AIR TEMPERATURE IN EUROPE

In this section we examine the influence of maritime-air advection in raising the winter air temperature in Europe. The strength of the warm advection is quantified by a specific Index, I_{na} , of the ocean-surface southwesterlies over the eastern North Atlantic. From the SSM/I dataset (Atlas et al. 1996), we evaluate at 45°N : 20°W pentad-averages of the wind speed from all the measurements which report the direction from the quadrant 180° – 270° . When the direction is not southwesterly, the speed is counted to the average as a zero speed. The Index I_{na} derived by this approach is plotted in Figure 1 for the winters 1989/90 and 1995/96, that is for the 24 pentads from December 2–6 to March 27–31.

Alongside I_{na} we plotted in Figure 1 the pentad-averaged T_{max} of daily maximum air temperature in Brussels, Belgium. We selected this Royal Meteorological Institute meteorological station, since its very long record establishes this dataset as suitable for the evaluation of trends (Demarée et al. 2001). We note very large differences between the air temperature T_{max} in 1989/90 and those in 1995/96, amounting to about 10°C for much of the winter. Based on detailed correlation analysis (Otterman et al. 1999), this difference in T_{max} can be attributed to the difference in the warm advection, quantified by I_{na} , the strength of the southwesterlies at the “gateway” to Europe. We note here that I_{na} exceeds 10 ms^{-1} for much of the warm 1989/90 period, but assumes low values, generally below 4 ms^{-1} , for most of the cold winter 1995/96. The differences in the warm advection affect the surface-air temperature over vast areas, as can be seen from Figure 2, where we compare February 1990 with February 1996. The observed sensitivity of the surface-air temperature to the low-level maritime-air advection in winter is directly due to the large air temperature difference prevailing then between the North Atlantic and the European continent (Otterman et al. 2000). The warming of the near-surface layer tends to produce a steeper lapse rate, that




Figure 1. Maximum daily air temperature (T_{\max}) in Brussels, and the Index of the southwesterlies (I_{na}), pentad-averages for the December-to-March periods in 1989/90 and 1995/96.

is, to destabilize the atmospheric conditions. These effects are analyzed in the next section.

COROLLARY EFFECTS OF WARM AND MOIST LOW-LEVEL ADVECTION

The winters of 1989/90 and 1995/96 are well representative of opposite extremes in the NAO index. Sea level pressure gradients across 25°W longitude between latitudes 45 and 55°N were approximately twice as strong in 1990 as in 1996. Some of the largest sea level pressure differences between the two NAO states occur typically around the Bay of Biscay (Rogers 1997). During periods such as early 1990, when storms migrate far northeastward of Iceland, they can advect warm air and cloud cover as far east as central Siberia (Rogers and Mosley-Thompson 1995).




Figure 2. Surface air temperature (T_s in $^{\circ}\text{C}$) over the eastern North Atlantic and mid-latitude Europe (46°N to 60°N ; 10°W to 50°E), February 1990 at the top, February 1996 in the middle, their difference in the lowest figure.

While only monthly (or seasonal) data are relevant in climate studies, insight into the phenomenology can be gained from a “single-moment” scenario. We illustrate the winter 1990 conditions in Figure 3 presenting surface winds (from SSM/I and ECMWF) for February 1, 1990, 00Z. We note a “STREAK” of southwesterlies directed toward France and England, where the wind speed exceeds 10 ms^{-1} .




Figure 3. Surface winds (ms^{-1}) over the North Atlantic and Europe, February 1, 1990, 00Z (from SSM/I and ECMWF).




Figure 4. Ascent rates at the 700 mb level (10 Pa s^{-1}), February 1, 1990, 12Z (from ECMWF).




Figure 5. As Figure 2, for the water vapor in the vertical column (kg/m^2).

The low-level warm advection in 1990 produced strong updrafts: ascent rates of up to -0.4 Pa s^{-1} were observed in monthly averages at 700 mb, which were especially strong over the ocean just to the west of Scandinavia. Such high monthly-average ascent rates persisted for more than a month. By comparison, in 1996, the monthly-average ascent rates at that level were reported generally as zero, with only occasional -0.1 Pa s^{-1} readings.




Figure 6. As Figure 2, for the total cloud cover (percent).

The way in which the strong warm advection illustrated in Figure 3 is affecting the state of the atmosphere 12 hours later, that is on February 1, 1990, 12Z is shown in Figure 4. We observe at 700 mb large cells of high ascent rates in a 270° “ring” around central Europe. In the strongest cells, one to the northwest of Ireland and the other centered on the southern coast of Scandinavia, the ascent rate tops -1.2 Pa s^{-1} . In cells over Finland, Spain and western Mediterranean, the ascent rates top -0.9 Pa s^{-1} .

The warm air masses advected in the 1989/90 winter are certainly moist, as we illustrate in Figure 5: in central Europe at latitudes 52-to-60°N the total precipitable water is some 5 kg m⁻² higher in February 1990 than in February 1996. Higher moisture levels and a steeper lapse rate combine to increase cloudiness: the total cloud cover reaches 60 to 70% for February 1990 as compared to 45 to 55% in 1996 (Fig. 6). In February the sun is low over the horizon in Europe, and the absorptivity (co-albedo) of the snow-covered surface is low, and thus absorption of insolation is hardly an important consideration for the surface temperature. Enhanced cloud cover and precipitable water vapour levels reduce heat loss to space. These corollary effects, by enhancing the greenhouse factor, reinforce the underline near-surface warming by the low-level warm advection from the North Atlantic.

DISCUSSION AND CONCLUSIONS

Characterizing climatic conditions for a region involves specifying average values of the key climatic parameters, combined with their variability, that is, the statistics of departures from the averages. Poleward of 35°, the key climate parameter for agriculture and forestry (and for heating-fuel demand) is the surface temperature (both of skin and the surface air, which are closely related). Surface air temperatures dictate the beginning of planting and seeding at the end of winter, and the harvesting of crops before the next winter sets in. The winter/early-spring differences between 1989/90 and 1995/1996 that we discuss here, persisting at the level of 10°C for more than a week at a time for Brussels (see Fig. 1), are extremely large. This reported variability is entirely consistent with the variations in the February means characterizing central Poland (see

Table 1. Air temperature (°C) in Warsaw, 1990 and 1996.

Monthly average air temperature (Lorenc 2000)		
Months	1990	1996
December, previous year	1.1	-4.8
January	1.9	-5.7
February	5.1	-5.3
March	6.7	-1.1
Monthly average maximum air temperature (absolute maximum air temperature in brackets) (Niedźwiedz and Ustrnul 1994)		
Months	1990	1996
December, previous year	3.7 (14.2)	-2.5 (7.7)
January	4.1 (10.7)	-3.2 (5.3)
February	9.1 (17.2)	-2.3 (5.3)
March	11.5 (20.5)	1.4 (6.4)

Table 2. Air temperature ($^{\circ}\text{C}$) in Toruń, 1990 and 1996.

Monthly average air temperature (highest and lowest mean daily air temperature in brackets)		
Months	1990	1996
December, previous year	1.2 (9.5/-5.7)	-5.0 (3.5/-14.1)
January	2.4 (7.8/-5.5)	-5.7 (2.1/-16.0)
February	5.3 (11.5/-0.1)	-5.7 (1.6/-15.6)
March	6.7 (11.7/0.4)	-1.0 (3.4/-4.9)
Monthly average maximum air temperature (absolute maximum air temperature in brackets)		
Months	1990	1996
December, previous year	3.7 (13.0)	-2.2 (6.5)
January	4.6 (12.0)	-3.3 (5.2)
February	9.3 (17.1)	-2.4 (4.8)
March	11.2 (21.5)	2.1 (7.8)

Table 1 or Table 4.1, for Warsaw p. 78, in Woś 1999). The fluctuations reported for Toruń, amount to 11.0°C and 11.7°C in the case of average monthly T_s and T_{\max} , respectively (see Table 2). The above-described large temperature differences occurred over almost the whole continental mid-latitude Europe (see Fig. 2).

Our study indicates that the underlying cause of the winter air temperature fluctuations in Europe is the variability of the surface winds over the North Atlantic. This dependence constitutes an important teleconnection similar to that examined by Hurrell (1996) and Otterman et al. (2000). The dependence of temperature, specifically in Poland, on the circulation patterns was discussed by Niedźwiedz and Ustrnul (1994). The strong winter temperature trends at northern latitudes in Eurasia reported by Ross et al. (1996) maybe the result of intensified advection from the North Atlantic. A more quantitative assessment of the sensitivity of the air temperatures in various European locations to the advection effects should be attempted.

REFERENCES

- Atlas R., Hoffman R. N., Bloom S. C., Jusem J. C., Ardizzone J., 1996, *A multiyear global surface wind velocity dataset using SSM/I wind observations*, Bull. Amer. Meteor. Soc., 77, 869–882.
- Demarée G., Lachaert P.-J., Verhoeve T., Thoen E., 2001, *The long-term daily Central Belgium Temperature (CBT) time-series 1767–2000. History of the Instrumental Meteorological Observations in Belgium*, invited paper, special issue, Climatic Change.
- Hurrell J. W., 1996, *Influence of variations in extratropical wintertime teleconnections in Northern Hemisphere temperature*, Geophys. Res. Lett., 23, 665–668.
- Jaagus J., Ahas R., 2000, *Space-time variations of climate seasons and their correlation with the phenological development of nature in Estonia*, Clim. Res. 15, 207–219.

- Kalnay E. et al., 1996, *The NCEP/NCAR 40-year reanalysis project*, Bull. Amer. Meteor. Soc., 77, 437–471.
- Kistler R. et al., 2001, *The NCEP/NCAR 50-year Reanalysis: Monthly-means CD-ROM and documentation*, Bull. Amer. Meteor. Soc., 82, 247–268.
- Lorenc H., 2000, *Studia nad 220-letnią (1779–1998) serią temperatury powietrza w Warszawie oraz ocena jej wiekowych tendencji (Studies of 220 years of air temperature dataset for Warsaw (1779–1998) and an estimation of secular tendencies)*, Materiały Badawcze IMGW, Seria: Meteorologia, 31, 104 pp. (in Polish).
- Menzel A., Fabian P., 1999, *Growing season extended in Europe*, Nature, 397, 65.
- Niedźwiedź T., Ustrnul Z., 1994, *Maximum and minimum temperatures in Poland and the variability of atmospheric circulation*, [in:] Brázdil R., Kolar M. (ed.), *Contemporary climatology*, Brno, 420–425.
- Otterman J., Atlas R., Ardizzone J., Starr D., Jusem J. C., Terry J., 1999, *Relationship of late-winter temperatures in Europe to North Atlantic surface winds: A correlation analysis*, Theor. and Appl. Climatol., 64, 201–211.
- Otterman J., Ardizzone J., Atlas R., Hu H., Jusem J. C., Starr D., 2000, *Winter-to-spring transition in Europe 48–54°N: from temperature control by advection to control by insolation*, Geophys. Res. Lett., 17, 561–564.
- Rogers J., 1997, *North Atlantic storm tract variability and its association to the North Atlantic oscillation and climate variability of Northern Europe*, J. Climate, 10, 1635–1647.
- Rogers J. C., Mosley-Thompson E., 1995, *Atlantic arctic cyclones and the mild Siberian winters of the 1980s*, Geophys. Res. Lett., 22, 799–802.
- Ross R. J., Otterman J., Starr D. O., Elliott W. P., Angell J. K., Susskind J., 1996, *Regional trends of surface and tropospheric temperature and evening-morning temperature difference in northern latitudes*, Geophys. Res. Lett., 23, 3179–3193.
- Woś A., 1999, *Klimat Polski (The climate of Poland)*, Wyd. Naukowe PWN SA, Warsaw, 302 pp.