

Flyunt Igor-Severyn S, Musiyenko Vira Y, Horkovenko Nataliya L, Fil' Vitaliy M, Ivassivka Anzhela S, Zukow Walery. Some hormonal, lipid and haemodynamic maintenances of alternative vegetotropic effects of balneotherapy on SPA Truskavets'. Journal of Education, Health and Sport. 2017;7(11):288-293. eISSN 2391-8306. DOI <http://dx.doi.org/10.5281/zenodo.1069598> <http://ojs.ukw.edu.pl/index.php/johs/article/view/5082>

The journal has had 7 points in Ministry of Science and Higher Education parametric evaluation. Part B item 1223 (26.01.2017).

1223 Journal of Education, Health and Sport eISSN 2391-8306 7

© The Authors 2017;

This article is published with open access at Licensee Open Journal Systems of Kazimierz Wielki University in Bydgoszcz, Poland

Open Access. This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited. This is an open access article licensed under the terms of the Creative Commons Attribution Non Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted, non commercial use, distribution and reproduction in any medium, provided the work is properly cited.

This is an open access article licensed under the terms of the Creative Commons Attribution Non Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted, non commercial use, distribution and reproduction in any medium, provided the work is properly cited.

The authors declare that there is no conflict of interests regarding the publication of this paper.

Received: 05.11.2017. Revised: 21.11.2017. Accepted: 30.11.2017.

SOME HORMONAL, LIPID AND HAEMODYNAMIC MAINTENANCES OF ALTERNATIVE VEGETOTROPIC EFFECTS OF BALNEOTHERAPY ON SPA TRUSKAVETS'

Igor-Severyn S Flyunt¹, Vira Y Musiyenko², Nataliya L Horkovenko³, Vitaliy M Fil'¹, Anzhela S Ivassivka¹, Walery Zukow⁴

¹Department of Anatomy, Physiology and Valeology, Ivan Franko Pedagogical University, Drohobych, Ukraine fillvitalij@gmail.com

²Clinical Sanatorium "Moldova", Truskavets', Ukraine

³JSC "Dnipro-Beskyd", Truskavets', Ukraine dni-pro-b@bk.ru

⁴Department of Spatial Management and Tourism, Faculty of Earth Sciences, Nicolaus Copernicus University, Torun, Poland w.zukow@wp.pl

Abstracts

Background. The data on the influence of balneotherapy in the spa Truskavets' on the autonomic nervous system are ambiguous, so research in this direction remains relevant. The purpose of this study is accompanying changes in some hormonal, lipid and hemodynamic parameters in women with alternative variants of the vegetotropic effect of balneotherapy. **Materials and Methods.** The object of the study were 30 women aged 30-50 years old, patients with chronic cholecystitis in the remission phase, combined with polycystic ovaries and hyperplasia of the thyroid gland. Before and after balneotherapy, HRV was recorded and hormonal, lipid as well as veloergometric parameters were determined. **Results.** We confirmed the ambivalent nature of the vegetotropic effect of balneotherapy: sympathicotonic in 57% women and vagotonic in 43%. It is revealed the tendency to decrease testosterone level with sympathicotonic effect and the opposite tendency with vagotonic effect. There were no significant differences with respect to thyroid hormones, FSH, estradiol, prolactin and cortisol. Vagotonic but not sympathicotonic effect is accompanied by a tendency to decrease the coefficient of atherogenicity and the response of systolic blood pressure to standard physical load. **Conclusion.** Alternative vegetotropic effects of balneotherapy on spa Truskavets' are accompanied by alternative changes in the level of testosterone alone, but no other registered hormones as well as lipid and hemodynamic parameters.

Keywords: spa Truskavets', balneotherapy, women, HRV, hormones, lipids, hemodynamics.

INTRODUCTION

Previously it was shown that the course use of bioactive water Naftussya influences the autonomic nervous system of healthy rats of both sexes [18,19,24], children of both sexes

with dysadaptoxis [17,19] and women with chronic endocrine-gynecological pathology [21,24-27] is ambiguous: it carries both vagotonic and sympathetic tonic effects, or does not significantly alter the vagal-sympathetic balance. Concomitant changes in some hormonal, metabolic and hemodynamic parameters are also ambiguous. On the other hand, the same variety has been found for thyrotropic effects of bioactive water Naftussya [2,4,5,9-16] as well as for changes in stress-index [23], hemodynamics, cholekinetics, gastric and pancreatic secretion [3]. So research in this direction remains relevant. The purpose of this study is accompanying changes in some hormonal, lipid and hemodynamic parameters in women with alternative variants of the vegetotropic effect of balneotherapy.

MATERIAL AND METHODS

The object of the study were 30 women aged 30-50 years old, patients with chronic cholecystitis in the remission phase, combined with polycystic ovaries who underwent rehabilitation course at the spa Truskavets'. At the arrival and after the completion of the three-week course of drinking bioactive water Naftusya (3,5 ml/kg for 1 hour before meals thrice a day), carried out vegetative, biochemical and veloergometric tests. The state of vegetative regulation is estimated by the HRV [1], using the hardware-software complex "CardioLab+HRV" (in the "KhAI-MEDICA", Kharkiv, Ukraine).

Biochemical tests included, firstly, the determination of the content of hormones in blood plasma: TSH, thyroxine, triiodothyronine, prolactin, FSH, estradiol, testosterone and cortisol (using the ELISA, the analyzer "Tecan", Oesterreich and the corresponding sets of reagents of Alkor Bio Co., Russian Federation [7]). Another set of tests related to the parameters of the lipid plasma spectrum: triglycerides and cholesterol of the lipoproteins of very low (VLD), low (LD) and high (HD) densities. The determination was carried out by means of reflectometry, using the apparatus "Reflotron" (BRD). Norms borrowed from literature [1,2,6,7,19,23,24,26].

Physical performance was estimated by the method of two-stage (load 0,5 and 1,5 W/kg) veloergometry. Veloergometer „Tunturi” (Finland) used.

The digital material is processed on a personal computer by the method of variation analysis using the program „Statistica 5.5”.

RESULTS AND DISCUSSION

By orientation of the dynamics of the sympathetic-vagal balance index (LF/HF), two groups of women subordinate to the vagotonic or sympathotonic vegetotropic effect of balneotherapy were retrospectively formed (Table 1).

Table 1. Features of hormonal status and its dynamics in alternative variants of vegetotropic effect of balneotherapy

Parameters	Term	Vegetotropic effect of balneotherapy		Norm
		Vagotonic (n=13)	Sympathotonic (n=17)	
LF/HF	Before	6,5±0,9*	2,2±0,4 ^v	1,5±0,1
	After	3,1±0,6* [#]	5,0±0,7* [#]	
TSH, mIU/L	Before	4,36±0,67*	6,21±0,77*	1,90±0,15
	After	4,81±0,74*	5,14±0,93*	
Thyroxine, nM/L	Before	96±10	93±10	110±4
	After	107±12	96±9	
Triiodothyronine, нМ/л	Before	1,79±0,26	1,63±0,30	2,10±0,09
	After	2,13±0,33	1,80±0,23	
Prolactin, μg/L	Before	15,9±2,9*	14,9±2,6*	8,4±0,5
	After	18,5±3,3*	16,3±2,8*	
FSH, IU/L	Before	5,7±0,5	5,5±0,4	6,1±0,4
	After	6,6±0,3	6,0±0,4	
Estradiol, ng/L	Before	94±8	97±9	115±8
	After	96±12	93±10	
Testosterone, μg/L	Before	0,80±0,14*	0,95±0,12*	0,28±0,02
	After	1,00±0,14*	0,80±0,16*	
Cortisol, μg/L	Before	198±10*	196±11*	165±8
	After	199±12*	194±11*	

Notes: 1. Parameters that are significantly different from normal are marked*.

2. Significant differences between the initial parameters of the groups are marked^v.

3. Significant differences between the endpoints and the starting points in each group are marked[#].

First, we will analyze the features of the initial neuro-endocrine status and the dynamics of its parameters. In the first place, it was found that the examined contingent is characterized by hypothyroidism as a whole, as evidenced by the lower levels of triiodothyronine combined with elevated levels of TSH. Hypothyroidism in women with normal or slightly elevated sympatho-vagal index is slightly more pronounced than in women with significant sympathicotonia: T₃ level are 78% of the mean norm (MN) versus 85% of MN, while TSH level, on the contrary, is 327% of MN against 229% of MN.

Hypothyroidism is associated with an equally moderate increase in cortisol levels (up to 119% and 120% of MN) and equally pronounced hyperprolactinemia (177% and 189% of MN), while the initial levels of follicle stimulating hormone and estradiol are almost normal. The listed hormones are not subject to essential influence of balneotherapy, though it should be noted the positive dynamics of thyroid hormones and FSH in the direction of normalization, in conjunction with the subsequent increase in hyperprolactinemia. However, these trends are not related to the nature of the vegetotropic effect.

Instead, the level of testosterone, which is significantly elevated, is somewhat higher in normotonics (339% of MN) than in sympathotonics (286% of MN), the opposite changes with respect to the dynamics of the sympatho-vagal index. In particular, for the sympathetic effect, it decreases (by 16%), whereas for vagotonic it increases (by 23%).

The lipid plasma spectrum (Table 2) as a whole in the contingent is characterized by severe hypertriglyceridemia: 148% of MN in the euthonics and 167% of MN in the sympatotonics, and an increase in cholesterol in the composition of very low density lipoproteins (149% and 168% of MN) and its decrease in composition lipoproteins low (up to 85% and 92% of MN) and high (up to 82% and 84% of MN, respectively) density. As a

result, it gives an equally pronounced increase in the Klimov's atherogenicity coefficient - up to 138% of MN.

The sympathetic effect of balneotherapy is associated with the absence of the dynamics of lipid profile parameters; instead, for vagotonic effect the tendency toward growth of very low density lipoproteins cholesterol is 17% in combination with the opposite trend of low density lipoproteins cholesterol (-21%), which, even in the absence of the dynamics of the antiatherogenic fraction, leads to small (by 13%) reduction of the atherogenicity.

Table 2. Features of lipid plasma spectrum and its dynamics in alternative variants of vegetotropic effect of balneotherapy

Parameters	Term	Vegetotropic effect of balneotherapy		Norm
		Vagotonic (n=13)	Sympathotonic (n=17)	
Triglycerides, mM/L	Before	2,10±0,16*	1,86±0,14*	1,26±0,04
	After	2,48±0,17*	1,90±0,15*	
Total cholesterol, mM/л	Before	5,00±0,14	4,68±0,11*	5,26±0,09
	After	4,49±0,11* [#]	4,64±0,11*	
VLDL cholesterol, mM/L	Before	0,69±0,05*	0,61±0,05*	0,41±0,01
	After	0,81±0,06*	0,61±0,05*	
LDL cholesterol, mM/L	Before	3,04±0,14	2,81±0,11*	3,32±0,07
	After	2,39±0,12* [#]	2,74±0,12*	
HDL cholesterol, mM/L	Before	1,28±0,04*	1,26±0,05*	1,53±0,02
	After	1,30±0,05*	1,29±0,05*	
Klimov's atherogenicity coefficient	Before	3,07±0,15*	3,06±0,16*	2,21±0,03
	After	2,66±0,13* [#]	2,86±0,14*	

The analysis of hemodynamic parameters (Table 3) shows that women subordinate to the vagotonic effect of balneotherapy are characterized by slightly higher initial levels of systolic and diastolic pressure at rest and after bicycle ergometric loading of 1,5 W/kg compared to women subordinate to sympathetic effect.

Table 3. Changes in hemodynamics in alternative variants of vegetotropic effect of balneotherapy

Parameters	Term	Vegetotropic effect of balneotherapy	
		Vagotonic (n=13)	Sympathotonic (n=17)
Systolic blood pressure at rest, mmHg	Before	125±4	119±4
	After	123±2	119±3
Diastolic blood pressure at rest, mmHg	Before	81±2	76±2 ^v
	After	79±1	76±1
Systolic blood pressure after loading, mmHg	Before	147±4	142±4
	After	135±3 [#]	140±4
Diastolic blood pressure after loading, mmHg	Before	88±2	79±2 ^v
	After	81±2 [#]	82±3
Heart rate after at rest, beats/min	Before	85,7±2,9	65,0±2,9 ^v
	After	76,8±3,1 [#]	75,6±4,1 [#]
Heart rate after loading, beats/min	Before	129±3	135±4
	After	138±4	133±3
Tachycardic-hypertensive reaction index, μW/kg•beat•mmHg	Before	77,7±4,2	78,4±4,1
	After	80,6±4,2	80,1±4,4

The difference in heart rate is especially noticeable at rest, but tachycardic responses to load are not significantly different. The sympathetic effect of balneotherapy is expected to be

accompanied by an increase in heart rate at rest at 16%, but the tachycardia response to exercise remains unchanged, as well as parameters of arterial pressure at rest and after loading. Instead, the vagotonic effect is associated with a 10% reduction in heart rate at rest, in combination with an increase of 7% after exercise, with a systolic pressure decrease of 8% and a diastolic decrease of 8% as well. The index of tachycardic-hypertensive response to a load of 1,5 W/kg, proposed by IL Popovych for the assessment of physical ability [3,4], remains unchanged in both groups.

CONCLUSION

Alternative vegetotropic effects of balneotherapy on spa Truskavets' are accompanied by alternative changes in the level of testosterone alone, but no other registered hormones as well as lipid and hemodynamic parameters.

ACCORDANCE TO ETHICS STANDARDS

This study was approved by the local ethical committee of Truskavets' Scientists Association. Tests in patients are conducted in accordance with positions of Helsinki Declaration 1975, revised and complemented in 2002, and directive of National Committee on ethics of scientific researches. During realization of tests from all participants the informed consent is got and used all measures for providing of anonymity of participants. For all authors any conflict of interests is absent.

REFERENCE

1. Baevskiy RM, Ivanov GG. Heart Rate Variability: theoretical aspects and possibilities of clinical application [in Russian]. *Ultrazvukovaya i funktsionalnaya diagnostika*. 2001; 3: 106-127.
2. Bul'ba AYa. Variants of the thyrotropic effects of balneotherapy in the spa Truskavets', their neuroendocrine and clinical satellites and predictors in women with thyroid hyperplasia [in Ukrainian]. *Medical Hydrology and Rehabilitation*. 2007; 5(2): 30-45.
3. Chebanenko OI, Chebanenko LO, Popovych IL. Multivariate Balneoeffects of Factors of Spa Truskavets' and Forecasting [in Ukrainian]. Kyiv: UNESCO-SOCIO. 2012: 496 p.
4. Fuchko OL. Hemodynamic accompaniment of the thyrotropic effects of balneotherapy in the spa Truskavets' [in Ukrainian]. *Medical Hydrology and Rehabilitation*. 2010; 8(2): 35-39.
5. Fuchko OL, Bul'ba AYa. Variants of the thyrotropic effects of balneotherapy in the spa Truskavets' in women with thyroid hyperplasia and associated changes in the parameters of lipid and electrolyte exchanges [in Ukrainian]. *Medical Hydrology and Rehabilitation*. 2008; 6(3): 51-59.
6. Goryachkovskiy AM. Clinical biochemistry [in Russian]. Odesa: Astroprint. 1998: 608 p.
7. Instructions for application for recruitment reagents for ELISA investigations hormones in the blood of humans [in Russian]. St. Petersburg: JSC "Alkor Bio". 2000.
8. Ivassivka SV, Bubnyak AB, Kovbasnyuk MM, Popovych IL. Genesis and role of phenols in waters from Naftussya layer [in Ukrainian]. In: *Problems of pathology in experiment and clinic. Scientific works of Drohobych Medical Institute*. Vol XV. Drohobych. 1994: 6-11.
9. Kozyavkina NV. Variants of thyrotropic effects of bioactive water Naftussya and their lipid support [in Ukrainian]. *Medical Hydrology and Rehabilitation*. 2008; 6(3): 115-122.
10. Kozyavkina NV. Neuro-endocrine and electrolyte accompaniment of polyvariant thyrotropic effects of bioactive water Naftussya [in Ukrainian]. *Medical Hydrology and Rehabilitation*. 2009; 7(1): 51-55.
11. Kozyavkina NV. Immune accompaniment of polyvariant thyrotropic effects of bioactive water Naftussya [in Ukrainian]. *Medical Hydrology and Rehabilitation*. 2009; 7(2): 27-39.
12. Kozyavkina NV. Polyvariance of thyrotropic effects of bioactive water Naftussya in women with chronic endocrine-gynecological pathology, their neuroendocrine-immune and clinical support and prediction capabilities [in Ukrainian]. *Medical Hydrology and Rehabilitation*. 2011; 9(2): 4-23.
13. Kozyavkina NV. Thyrotropic effects of bioactive water Naftussya in female rats and their

metabolic, neuroendocrine and immune accompaniments [in Ukrainian]. *Medical Hydrology and Rehabilitation*. 2012; 10(4): 91-113.

14. Kozyavkina NV, Barylyak LG, Yanchij OR, Fuchko OL Thyrotropic effects of Naftussya water, their vegetative manifestation and the possibility of prediction [in Ukrainian]. *Fiziologichnyi Zhurnal*. 2013; 59(6): 81-87.

15. Kozyavkina NV, Gozhenko AI, Barylyak LG, Korolyshyn TA, Popovych IL. Polyvariance of immediate thyrotropic effects of bioactive water Naftussya, their neuroendocrine-immune support and the possibility of prediction [in Ukrainian]. *Medical Hydrology and Rehabilitation*. 2013; 11(4): 27-54.

16. Kozyavkina NV, Popovych IL, Zukow W. Metabolic accompaniment of thyrotropic effects of bioactive water Naftussya at the women with thyroid hyperplasia. *Journal of Health Sciences*. 2013; 3(5): 409-424.

17. Kozyavkina OV. Vegetotropic effects of bioactive water Naftussya at children with dysfunction of the neuroendocrine-immune complex, these endocrine-immune accompaniment and the ability to predict [in Ukrainian]. *Medical Hydrology and Rehabilitation*. 2011; 9(2): 24-39.

18. Kozyavkina OV. Vegetotropic effects of bioactive water Naftussya at males rats and their endocrine, electrolyte and immune accompaniments [in Ukrainian]. *Medical Hydrology and Rehabilitation*. 2012; 10(3): 65-92.

19. Kozyavkina OV, Kozyavkina NV, Gozhenko OA, Gozhenko AI, Barylyak LG, Popovych IL. Bioactive Water Naftussya and Neuro-endocrine-immune Complex [in Ukrainian]. Kyiv: UNESCO-SOCIO. 2015: 349 p.

20. Kozyavkina OV, Popovych IL, Zukow W. Immediate vegetotropic effects of bioactive water Naftussya and their neuro-endocrine-immune accompaniment in healthy men. *Journal of Health Sciences*. 2013; 3(5): 391-408.

21. Kozyavkina OV, Vis'tak HI, Popovych IL. Factor, canonical and discriminant analysis of vegetotropic effects and accompanying changes for thyroide, metabolic and haemodynamic parameters at the women, caused by bioactive water Naftussya [in Ukrainian]. *Medical Hydrology and Rehabilitation*. 2013; 11(3): 4-28.

22. Popovych IL. The concept of neuro-endocrine-immune complex (review) [in Russian]. *Medical Hydrology and Rehabilitation*. 2009; 7(3): 9-18.

23. Popovych IL, Barylyak LG. Influence of course of use of bioactive water Naftussya on the level of stress in women with endocrine-gynecological pathology [in Ukrainian]. *Medical Hydrology and Rehabilitation*. 2009; 7(3): 100-118.

24. Popovych IL, Vis'tak HI, Gumega MD, Ruzhylo SV. Vegetotropic Effects of Bioactive Water Naftussya and their Endocrine-Immune, Metabolic and Hemodynamic Accompaniments [in Ukrainian]. Kyiv: UNESCO-SOCIO. 2014: 163 p.

25. Vis'tak HI. Endocrine and immune support of polyvariant vegetotonic effects of bioactive water Naftussya in women [in Ukrainian]. *Medical Hydrology and Rehabilitation*. 2009; 7(3): 81-85.

26. Vis'tak HI. Interrelationships between vegetotropic and endocrine, immunotropic as well as clinical effects of bioactive water Naftussya in women with hyperplasia of the thyroid gland [in Ukrainian]. *Medical Hydrology and Rehabilitation*. 2012; 10(2): 37-66.

27. Vis'tak HI, Kozyavkina OV, Popovych IL, Zukow W. Vegetotropic effects of bioactive water Naftussya spa Truskavets' and their thyroide, metabolic and haemodynamic accompaniments at the women. *Journal of Health Sciences*. 2013; 3(10): 557-582.