

Kwiatkowska Klaudia, Lamtych Martyna, Kubiak Karolina, Litwa Ewelina, Badiuk Nataliia. Assessment of fatigue and autonomic dysfunction in myasthenia gravis – case report. *Journal of Education, Health and Sport*. 2019;9(5):667-675. eISSN 2391-8306. DOI <http://dx.doi.org/10.5281/zenodo.3375836>
<http://ojs.ukw.edu.pl/index.php/johs/article/view/7322>

The journal has had 7 points in Ministry of Science and Higher Education parametric evaluation. Part b item 1223 (26/01/2017).
1223 Journal of Education, Health and Sport eISSN 2391-8306 7

© The Author(s) 2019;

This article is published with open access at Licensee Open Journal Systems of Kazimierz Wielki University in Bydgoszcz, Poland
Open Access. This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author (s) and source are credited. This is an open access article licensed under the terms of the Creative Commons Attribution Non commercial license (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted, non commercial use, distribution and reproduction in any medium, provided the work is properly cited.
This is an open access article licensed under the terms of the Creative Commons Attribution Non commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted, non commercial use, distribution and reproduction in any medium, provided the work is properly cited.
The authors declare that there is no conflict of interests regarding the publication of this paper.
Received: 19.05.2019. Revised: 22.05.2019. Accepted: 31.05.2019.

Assessment of fatigue and autonomic dysfunction in myasthenia gravis – case report

Klaudia Kwiatkowska¹, Martyna Lamtych¹, Karolina Kubiak¹,
Ewelina Litwa², Nataliia Badiuk³

¹Student Scientific Circle at Chair of Hygiene, Epidemiology and Ergonomics, Faculty of Health Sciences, Nicolaus Copernicus University in Toruń, Collegium Medicum in Bydgoszcz, Poland

²Scientific Circle at Department of Hygiene, Epidemiology and Ergonomics. Division of Ergonomics and Exercise Physiology, Nicolaus Copernicus University in Toruń, Collegium Medicum in Bydgoszcz, Poland

³State Enterprise Ukrainian Research Institute for Medicine of Transport, Ministry of Health of Ukraine, Odesa, Ukraine

Abstract

Introduction: Myasthenia gravis (MG) is an autoimmune disease in which neuromuscular conduction disorders occur. The reason is the loss of acetylcholine receptors and the presence of antibodies directed against these receptors. The most characteristic symptoms of myasthenia gravis are: tenderness and weakness of striated muscles, which intensify particularly during exercise and in the evening. Non-motor symptoms such as: fatigue,

disorders of the autonomic nervous system can be asymptomatic, and in a significant proportion of cases significantly affect the quality of life. A full understanding of non-motor symptoms is needed to treat patients with MG. The aim of the study was to present the characteristics of non-motor symptoms on the example of a patient with myasthenia gravis.

Material and methods: A 50-year-old woman who had been suffering from myasthenia gravis for 23 years was examined. Chalder Fatigue Scale, Epworth Sleepiness Scale, Orthostatic Grading Scale were used to evaluate non-motor symptoms. The orthostatic reaction was tested using the tilt test.

Results: The most frequently reported non-motor symptoms in the patient were: chronic fatigue, sleep disturbances and cardiovascular symptoms (orthostatic intolerance, weakness).

Conclusions: Myasthenia gravis is characterized by a large variability of symptoms. An important part of the clinical picture are non-motor symptoms that can affect the functioning of patients in everyday life.

Key words: myasthenia gravis, non-motor symptoms, fatigue

Introduction

Myasthenia gravis (MG) is one of the autoimmune diseases in which neuromuscular transmission disorders occur. The disease is in many cases late diagnosed by a heterogeneous clinical picture. The most characteristic symptoms for MG are excessive fatigue and muscle weakness, however, many patients may have additional non-motor symptoms such as fatigue or disorders of the autonomic nervous system [1, 2].

Fatigue is a subjective symptom accompanied by a feeling of general weakness. Occurs as an additional symptom of neurological (Parkinson's disease, Multiple Sclerosis), cardiovascular or neuromuscular diseases. Fatigue in patients with myasthenia gravis has not been as well understood as in other diseases. The reason may be that it is often confused with muscular fatigue, which is very characteristic for MG [3, 4].

Similar to fatigue, autonomic dysfunction has been well known in many other disease entities - including in multiple sclerosis, Parkinson's disease, Guillain-Barry syndrome, in cerebrovascular disease. In the available publications, the authors notice that symptoms such as heat intolerance, sweating disorders, orthostatic hypotension, urinary and sexual disorders, excessive dryness or diarrhea and constipation are also present in many patients with myasthenia gravis. However, these are not symptoms characteristic of MG, they are often asymptomatic. They are difficult to diagnose [2, 5, 6].

Case report

A woman aged 50, suffering from myasthenia gravis for 23 years. Currently unemployed, currently on a pension. In the interview, the patient reported arthralgia, progressive muscular weakness of the lower limbs and excessive fatigue, which is intensified under the influence of physical activity. A patient walking alone, without orthopedic assistance. Additional overweight (BMI - 27.7; height - 159 cm; weight - 70 kg).

In the assessment of autonomic disorders, the patient reported such symptoms as: orthostatic disorders, arrhythmia, dizziness, thermoregulation disorders, dry eyes and mouth, sleep disorders and secretory disorders (Table 1).

Table 1. Autonomic dysfunction in a patient.

Orthostatic disorders	+
Arrhythmia episodes	+
Vertigo	+
Constipation	-
Dry eyes and mouth	+
Secretory disorders	+
Stomach ache	-
Sleep disorders	+
Post-meal symptoms	-
Anxiety disorders	+
Diarrhoea	-
Urinary bladder dysfunctions	-
Sexual dysfunctions	-
Thermoregulatory disorders	+

Assessment of fatigue and autonomic nervous system functions

Chalder Fatigue Scale (CFS) was used to assess fatigue. It is a questionnaire consisting of 11 questions, of which the first seven are related to physical fatigue, the last seven are about mental fatigue [7]. The Epworth Sleepiness Scale (ESS) was used to assess sleepiness. This scale assesses the possibility of falling asleep in 8 everyday situations on a scale of 0 (impossible to sleep) to 3 (high probability of falling asleep). The higher the result achieved, the greater the tendency to fall asleep in everyday life [8].

The orthostatic reaction was tested by tilt test using Task Force Monitor® (TFM). It is a non-invasive method by which the autonomic and cardiovascular systems can be assessed. TFM consists of the following elements [9, 10]:

- impedance cardiography (IKG),
- pulse oximeter
- a device for continuous measurement of blood pressure,
- electrocardiograph,
- device for oscillometric blood pressure measurement.

The verticalization to 70 degrees took 5.20 minutes. The following parameters were analyzed: heart rate (HR), systolic blood pressure (sBP), diastolic blood pressure (dBP) and

mean blood pressure (mBP). The patient was informed about the methodology of the study, which she agreed to.

Additionally, the 5-point OGS (Orthostatic Grading Scale) questionnaire was used to assess the symptoms of orthostatic intolerance. The result obtained by the respondent is proportional to orthostatic intolerance [11].

Results

The body's physiological response to passive standing is to increase the heart rate (by up to 30%). Mean blood pressure does not change much [5]. The patient had an increase in the parameters studied, however, the increase in heart rate was very small, which may be manifested in the patient by dizziness or orthostatic intolerance (Fig. 1).




Figure 1. Hemodynamic values before and during the tilt test.

The results obtained in the tilt test are confirmed by the interview with the subject, who reported symptoms from the autonomic system - including orthostatic disorders, dizziness and arrhythmia (table 1).

The results of the questionnaires along with the interpretation are presented in Table 2. In CFS, the patient scored 16 points, which indicates excessive fatigue and weakness. ESS analysis does not indicate excessive sleepiness (4 points obtained). In addition, in the

Orthostatic Grading Scale, assessing the symptoms of orthostatic intolerance, the subject obtained 12 points out of 20 possible.

Table 2. Assessment of non-motor symptoms using OGS, CFS, ESS.

Scale	Score	Interpretation
Orthostatic Grading Scale (OGS)	12	<ul style="list-style-type: none"> • the higher the score, the greater the orthostatic intolerance; • maximum number of points: 20.
Chalder Fatigue Scale (CFS)	16	A score above 9 points is a fatigue.
Epworth Sleepiness Scale (ESS)	4	A score of 0-5 is no excessive sleepiness.

Discussion

The test organism's response to the tilt test, the results obtained in individual questionnaires and the interview signalize a reduced of the orthostatic reaction and may signal disorders of the autonomic nervous system (orthostatic disorders, arrhythmia, dizziness, dry eyes and mouth, secretory disorders, sleep disorders , anxiety disorders and thermoregulation disorders) and excessive fatigue. Currently, there are more and more scientific studies in which the authors raise the issue of fatigue and other non-motor symptoms in myasthenia gravis. Clinicians note that these are significant problems in this group of patients, negatively affecting the patient's quality of life [12-16].

In a prospective study, Nikolić et al. [12] compared the incidence of autonomic disorders due to the presence of antibodies and thymus disorders. In the group of patients with anti-acetylcholine receptor (anti-AChR) antibodies and thymoma, symptoms of autonomic dysfunction were in 86% of patients, with the most common occurrence: heat intolerance (76%), urinary bladder dysfunctions (81%) and sexual disorders (67%). Autonomic disorders in patients with anti-AChR positive antibodies, but without thymoma - were in 57% of

patients. In turn, in patients with anti-MuSK (anti-tyrosine kinase antibodies) the most common disorders were dizziness (39%), heat intolerance (30%) as well as dry mouth and sexual dysfunction - present in 26% of respondents. In the study by Elsais et al. [13], the most frequently reported symptoms of autonomic dysfunction in myasthenia gravis were excessive dryness (50%) and sleep disturbance (46%) and orthostatic disorder (44%). 32% reported vascular disorders, while food disorders - 22%.

Hoffmann et al. [14] used Chalder Fatigue Scale to assess fatigue. For 110 people with myasthenia gravis participating in the study, excessive fatigue was found in 56.1% of respondents. Elsais et al. [13] based on the same questionnaire noted the occurrence of fatigue in 44% of respondents. Alekseeva et al. [15] examined 73 people with myasthenia gravis using the FSS (Fatigue Severity Scale) questionnaire. Fatigue was found in up to 70% of respondents. In another study, Paul et al. [16] observed fatigue in 82% of patients with MG.

Non-motor symptoms, including disorders of the autonomic nervous system, are a big problem for patients with myasthenia gravis. These are not characteristic symptoms of myasthenia gravis, they are often asymptomatic and therefore their diagnosis may be a problem.

Conclusions

Myasthenia gravis is a disease characterized by a large variability and variety of symptoms. An important part of the clinical picture are non-motor symptoms such as chronic fatigue, sleep disturbances, orthostatic intolerance, weakness or thermoregulation disorders. They can affect the functioning of patients in everyday life.

All symptoms should be considered when diagnosing myasthenia gravis and when planning comprehensive therapeutic management.

References

1. Kostera-Pruszczyk A: Postępowanie terapeutyczne w miastonii. *Polski Przegląd Neurologiczny* 2013; 9 (1): 11–15.
2. Hamed SA: Comorbid nervous system manifestations and disorders with myasthenia gravis: Evidences and possible mechanisms. *Journal of Neurology and Neuroscience*. 2012; 3: 1:3.

3. Broła W, Ziomek M, Czernicki J: Fatigue syndrome in chronic neurological disorders. *Neurologia i Neurochirurgia Polska* 2007; 41 (4): 340–349.
4. Alekseeva TM, Gavrilov YV, Kreis OA, Valko PO, Weber KP, Valko Y: Fatigue in patients with myasthenia gravis. *Journal of Neurology* 2018; 265: 2312–2321.
5. Jörg J. Diagnostyka autonomicznego układu nerwowego i zaburzeń snu. *Urban & Partner, Wrocław* 2001.
6. Tong O, Delfiner L, Herskovitz S: Pain, Headache, and Other Non-motor Symptoms in Myasthenia Gravis. *Curr. Pain Headache Rep.* 2018; 22: 39.
7. Jackson C: The Chalder Fatigue Scale (CFQ 11). *Occupational Medicine.* 2015; 65 (1): 86.
8. Johns MW: A New Method for Measuring Daytime Sleepiness: The Epworth Sleepiness Scale. *Sleep* 1991; 14 (6): 540-545.
9. Zalewski P, Słomiński K, Klawe J, Tafil-Klawe M: Functional assessment of autonomic nervous system by using the Task Force Monitor® system. *Acta Bio-Optica Inf. Med. Inż. Biomed.* 2008; 14 (3): 228-234.
10. Zalewski P, Klawe J, Pawlak J, Tafil-Klawe M, Bitner A, Lewandowski A: Changes of the heart muscle contractility in healthy persons after single whole-body cryostimulation procedure. *Acta Bio-Optica Inf. Med. Inż. Biomed.* 2011; 17 (4): 257-261.
11. Schrezenmaier C, Gehrking JA, Hines SM, Low PA, Benrud-Larson LM, Sandroni P: Evaluation of orthostatic hypotension: relationship of a new self-report instrument to laboratory-based measures. *Mayo Clin Proc.* 2005; 80 (3): 330-334.
12. Nikolić A, Perić S, Nisić T, Popović S, Ilić M, Rakocević Stojanović V, Lavrnić D: The presence of dysautonomia in different subgroups of myasthenia gravis patients. *J Neurol.* 2014; 261: 2119–2127.
13. Elsaís A, Wyller VB, Håvard Loge J, Kerty E: Fatigue in myasthenia gravis: is it more than muscular weakness? *BMC Neurol.* 2013; 13: 132.
14. Hoffmann S, Ramm J, Grittner U, Kohler S, Siedler J, Meisel A: Fatigue in myasthenia gravis: risk factors and impact on quality of life. *Brain Behav.* 2016; 6 (10): e00538.
15. Alekseeva TM, Gavrilov YV, Kreis OA, Valko PO, Weber KP, Valko Y: Fatigue in patients with myasthenia gravis. *Journal of Neurology.* 2018; 265: 2312–2321.

16. Paul RH, Cohen RA, Goldstein JM, Gilchrist JM: Fatigue and its impact on patients with myasthenia gravis. *Muscle Nerve*. 2000; 23 (9): 1402-1406.

17.